COX IV Rabbit mAb
Catalog # AP78656

Specification

COX IV Rabbit mAb - Product Information

Application	**WB, IHC, IF**
Primary Accession	**P13073**
Reactivity	**Human, Mouse**
Host	**Rabbit**
Clonality	**Monoclonal Antibody**
Calculated MW	**19577**

COX IV Rabbit mAb - Additional Information

Gene ID 1327

Other Names
COX4I1

Dilution
WB~~1/500-1/1000
IHC~~1/100-1/200
IF~~1/50-1/200

Format
Liquid

COX IV Rabbit mAb - Protein Information

Name COX4I1 (HGNC:2265)

Function
Component of the cytochrome c oxidase, the last enzyme in the mitochondrial electron transport chain which drives oxidative phosphorylation. The respiratory chain contains 3 multisubunit complexes succinate dehydrogenase (complex II, CII), ubiquinol- cytochrome c oxidoreductase (cytochrome b-c1 complex, complex III, CIII) and cytochrome c oxidase (complex IV, CIV), that cooperate to transfer electrons derived from NADH and succinate to molecular oxygen, creating an electrochemical gradient over the inner membrane that drives transmembrane transport and the ATP synthase. Cytochrome c oxidase is the component of the respiratory chain that catalyzes the reduction of oxygen to water. Electrons originating from reduced cytochrome c in the intermembrane space (IMS) are transferred via the dinuclear copper A center (CU(A)) of subunit 2 and heme A of subunit 1 to the active site in subunit 1, a binuclear center (BNC) formed by heme A3 and copper B (CU(B)). The BNC reduces molecular oxygen to 2 water molecules using 4 electrons from cytochrome c in the IMS and 4 protons from the mitochondrial matrix.

Cellular Location
Mitochondrion inner membrane; Single-pass membrane protein

Tissue Location
Ubiquitous.

COX IV Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

COX IV Rabbit mAb - Images

abcepta

COX IV Antibody