

**Stathmin 1 Rabbit mAb**  
Catalog # AP78662**Specification**

---

**Stathmin 1 Rabbit mAb - Product Information**

Application	WB, IHC
Primary Accession	<a href="#">P16949</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Monoclonal Antibody
Calculated MW	17303

**Stathmin 1 Rabbit mAb - Additional Information****Gene ID** 3925**Other Names**  
STMN1**Dilution**  
WB~~1/500-1/1000  
IHC~~1/100-1/200**Format**  
Liquid**Stathmin 1 Rabbit mAb - Protein Information****Name** STMN1**Synonyms** C1orf215, LAP18, OP18**Function**  
Involved in the regulation of the microtubule (MT) filament system by destabilizing microtubules. Prevents assembly and promotes disassembly of microtubules. Phosphorylation at Ser-16 may be required for axon formation during neurogenesis. Involved in the control of the learned and innate fear (By similarity).**Cellular Location**  
Cytoplasm, cytoskeleton.**Tissue Location**  
Ubiquitous. Expression is strongest in fetal and adult brain, spinal cord, and cerebellum, followed by thymus, bone marrow, testis, and fetal liver. Expression is intermediate in colon, ovary, placenta, uterus, and trachea, and is readily detected at substantially lower levels in all other tissues examined. Lowest expression is found in adult liver. Present in much greater abundance in cells from patients with acute leukemia of different subtypes than in normal peripheral blood lymphocytes, non-leukemic proliferating lymphoid cells, bone marrow cells, or cells from patients

with chronic lymphoid or myeloid leukemia.

### Stathmin 1 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Stathmin 1 Rabbit mAb - Images

