

**DPP3 Rabbit mAb**  
Catalog # AP78787**Specification**

---

**DPP3 Rabbit mAb - Product Information**

Application	<b>WB</b>
Primary Accession	<a href="#">O9NY33</a>
Reactivity	<b>Human</b>
Host	<b>Rabbit</b>
Clonality	<b>Monoclonal Antibody</b>
Calculated MW	<b>82589</b>

**DPP3 Rabbit mAb - Additional Information****Gene ID** 10072**Other Names**

DPP3

**Dilution**

WB~~1/500-1/1000

**Format**

Liquid

**DPP3 Rabbit mAb - Protein Information****Name** DPP3**Function**

Cleaves and degrades bioactive peptides, including angiotensin, Leu-enkephalin and Met-enkephalin (PubMed:<a href="http://www.uniprot.org/citations/1515063" target="\_blank">1515063</a>, PubMed:<a href="http://www.uniprot.org/citations/3233187" target="\_blank">3233187</a>). Also cleaves Arg-Arg-beta-naphthylamide (in vitro) (PubMed:<a href="http://www.uniprot.org/citations/11209758" target="\_blank">11209758</a>, PubMed:<a href="http://www.uniprot.org/citations/3233187" target="\_blank">3233187</a>, PubMed:<a href="http://www.uniprot.org/citations/9425109" target="\_blank">9425109</a>).

**Cellular Location**

Cytoplasm, cytosol

**Tissue Location**

Detected in placenta (at protein level) (PubMed:3233187). Detected in erythrocytes (at protein level) (PubMed:1515063).

**DPP3 Rabbit mAb - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **DPP3 Rabbit mAb - Images**

