

CYP2C9 Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP7881c

Specification

CYP2C9 Antibody (Center) - Product Information

Application	WB, IHC-P,E
Primary Accession	P11712
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	235-265

CYP2C9 Antibody (Center) - Additional Information

Gene ID 1559

Other Names

Cytochrome P450 2C9, 11413-, (R)-limonene 6-monooxygenase, (S)-limonene 6-monooxygenase, (S)-limonene 7-monooxygenase, CYP11C9, Cytochrome P-450MP, Cytochrome P450 MP-4, Cytochrome P450 MP-8, Cytochrome P450 PB-1, S-mephenytoin 4-hydroxylase, CYP2C9, CYP2C10

Target/Specificity

This CYP2C9 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 235-265 amino acids from the Central region of human CYP2C9.

Dilution

WB~~1:1000
IHC-P~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CYP2C9 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

CYP2C9 Antibody (Center) - Protein Information

Name CYP2C9 {ECO:0000303|PubMed:11950794, ECO:0000312|HGNC:HGNC:2623}

Function A cytochrome P450 monooxygenase involved in the metabolism of various endogenous

substrates, including fatty acids and steroids (PubMed:[12865317](#), PubMed:[15766564](#), PubMed:[19965576](#), PubMed:[21576599](#), PubMed:[7574697](#), PubMed:[9435160](#), PubMed:[9866708](#)). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (NADPH--hemoprotein reductase) (PubMed:[12865317](#), PubMed:[15766564](#), PubMed:[19965576](#), PubMed:[21576599](#), PubMed:[7574697](#), PubMed:[9435160](#), PubMed:[9866708](#)). Catalyzes the epoxidation of double bonds of polyunsaturated fatty acids (PUFA) (PubMed:[15766564](#), PubMed:[19965576](#), PubMed:[7574697](#), PubMed:[9866708](#)). Catalyzes the hydroxylation of carbon-hydrogen bonds. Metabolizes cholesterol toward 25-hydroxycholesterol, a physiological regulator of cellular cholesterol homeostasis (PubMed:[21576599](#)). Exhibits low catalytic activity for the formation of catechol estrogens from 17beta- estradiol (E2) and estrone (E1), namely 2-hydroxy E1 and E2 (PubMed:[12865317](#)). Catalyzes bisallylic hydroxylation and hydroxylation with double-bond migration of polyunsaturated fatty acids (PUFA) (PubMed:[9435160](#), PubMed:[9866708](#)). Also metabolizes plant monoterpenes such as limonene. Oxygenates (R)- and (S)-limonene to produce carveol and perillyl alcohol (PubMed:[11950794](#)). Contributes to the wide pharmacokinetics variability of the metabolism of drugs such as S-warfarin, diclofenac, phenytoin, tolbutamide and losartan (PubMed:[25994031](#)).

Cellular Location


Endoplasmic reticulum membrane; Peripheral membrane protein. Microsome membrane; Peripheral membrane protein

CYP2C9 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CYP2C9 Antibody (Center) - Images



Western blot analysis of anti-CYP2C9 Antibody (Center)(Cat.#AP7881c) in CEM cell line lysates (35ug/lane).CYP2C9(arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human hepatocarcinoma tissue reacted with CYP2C9 antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

CYP2C9 Antibody (Center) - Background

CYP2C9 is a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. The protein localizes to the endoplasmic reticulum and its expression is induced by rifampin. The enzyme is known to metabolize many xenobiotics, including phenytoin, tolbutamide, ibuprofen and S-warfarin.

CYP2C9 Antibody (Center) - References

Nelson,D.R., Pharmacogenetics 14 (1), 1-18 (2004)

CYP2C9 Antibody (Center) - Citations

- [Comparative Analysis of Immunohistochemical Staining Intensity Determined by Light Microscopy, ImageJ and QuPath in Placental Hofbauer Cells](#)
- [Time-dependent expression pattern of cytochrome P450 epoxygenases and soluble epoxide hydrolase in normal human placenta.](#)