

**RNF7 Rabbit mAb**  
Catalog # AP78829**Specification****RNF7 Rabbit mAb - Product Information**

Application	<b>WB</b>
Primary Accession	<a href="#">Q9UBF6</a>
Reactivity	<b>Human, Mouse, Rat</b>
Host	<b>Rabbit</b>
Clonality	<b>Monoclonal Antibody</b>
Calculated MW	<b>12683</b>

**RNF7 Rabbit mAb - Additional Information**

Gene ID 9616

**Other Names**

RNF7

**Dilution**

WB~~1/500-1/1000

**Format**

Liquid

**RNF7 Rabbit mAb - Protein Information**Name RNF7 ([HGNC:10070](#))**Function**

Catalytic component of multiple cullin-5-RING E3 ubiquitin- protein ligase complexes (ECS complexes), which mediate the ubiquitination and subsequent proteasomal degradation of target proteins (PubMed: [21980433](http://www.uniprot.org/citations/21980433), PubMed: [33268465](http://www.uniprot.org/citations/33268465), PubMed: [38418882](http://www.uniprot.org/citations/38418882), PubMed: [38574733](http://www.uniprot.org/citations/38574733)). It is thereby involved in various biological processes, such as cell cycle progression, signal transduction and transcription (PubMed: [21980433](http://www.uniprot.org/citations/21980433), PubMed: [33268465](http://www.uniprot.org/citations/33268465), PubMed: [38418882](http://www.uniprot.org/citations/38418882), PubMed: [38574733](http://www.uniprot.org/citations/38574733)). The functional specificity of the E3 ubiquitin- protein ligase ECS complexes depend on the variable SOCS box-containing substrate recognition component (PubMed: [21980433](http://www.uniprot.org/citations/21980433), PubMed: [33268465](http://www.uniprot.org/citations/33268465)). Within ECS complexes, RNF7/RBX2 recruits the E2 ubiquitination enzyme to the complex via its RING-type and brings it into close proximity to the substrate (PubMed: [21980433](http://www.uniprot.org/citations/21980433), PubMed: [33268465](http://www.uniprot.org/citations/33268465)).

<http://www.uniprot.org/citations/34518685> target="\_blank">34518685</a>). Catalytic subunit of various SOCS- containing ECS complexes, such as the ECS(SOCS7) complex, that regulate reelin signaling by mediating ubiquitination and degradation of DAB1 (By similarity). The ECS(SOCS2) complex mediates the ubiquitination and subsequent proteasomal degradation of phosphorylated EPOR and GHR (PubMed:<a href="http://www.uniprot.org/citations/21980433" target="\_blank">21980433</a>, PubMed:<a href="http://www.uniprot.org/citations/25505247" target="\_blank">25505247</a>). Promotes ubiquitination and degradation of NF1, thereby regulating Ras protein signal transduction (By similarity). As part of the ECS(ASB9) complex, catalyzes ubiquitination and degradation of CKB (PubMed:<a href="http://www.uniprot.org/citations/33268465" target="\_blank">33268465</a>). The ECS(SPSB3) complex catalyzes ubiquitination of nuclear CGAS (PubMed:<a href="http://www.uniprot.org/citations/38418882" target="\_blank">38418882</a>). As part of some ECS complex, catalyzes 'Lys-11'-linked ubiquitination and degradation of BTRC (PubMed:<a href="http://www.uniprot.org/citations/27910872" target="\_blank">27910872</a>). ECS complexes and ARIH2 collaborate in tandem to mediate ubiquitination of target proteins; ARIH2 mediating addition of the first ubiquitin on CRLs targets (PubMed:<a href="http://www.uniprot.org/citations/34518685" target="\_blank">34518685</a>, PubMed:<a href="http://www.uniprot.org/citations/38418882" target="\_blank">38418882</a>). Specifically catalyzes the neddylation of CUL5 via its interaction with UBE2F (PubMed:<a href="http://www.uniprot.org/citations/19250909" target="\_blank">19250909</a>). Does not catalyze neddylation of other cullins (CUL1, CUL2, CUL3, CUL4A or CUL4B) (PubMed:<a href="http://www.uniprot.org/citations/19250909" target="\_blank">19250909</a>). May play a role in protecting cells from apoptosis induced by redox agents (PubMed:<a href="http://www.uniprot.org/citations/10082581" target="\_blank">10082581</a>).

**Cellular Location**

Cytoplasm. Nucleus

**Tissue Location**

Expressed in heart, liver, skeletal muscle and pancreas. At very low levels expressed in brain, placenta and lung

**RNF7 Rabbit mAb - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**RNF7 Rabbit mAb - Images**

