

**NR2C2/TR4 Rabbit mAb**  
Catalog # AP78856**Specification**

---

**NR2C2/TR4 Rabbit mAb - Product Information**

Application	<b>WB</b>
Primary Accession	<a href="#">P49116</a>
Reactivity	<b>Human, Mouse, Rat</b>
Host	<b>Rabbit</b>
Clonality	<b>Monoclonal Antibody</b>
Calculated MW	<b>65414</b>

**NR2C2/TR4 Rabbit mAb - Additional Information****Gene ID** 7182**Other Names**  
NR2C2**Dilution**  
WB~~1/500-1/1000**Format**  
Liquid**NR2C2/TR4 Rabbit mAb - Protein Information****Name** NR2C2**Synonyms** TAK1, TR4**Function**

Orphan nuclear receptor that can act as a repressor or activator of transcription. An important repressor of nuclear receptor signaling pathways such as retinoic acid receptor, retinoid X, vitamin D3 receptor, thyroid hormone receptor and estrogen receptor pathways. May regulate gene expression during the late phase of spermatogenesis. Together with NR2C1, forms the core of the DRED (direct repeat erythroid-definitive) complex that represses embryonic and fetal globin transcription including that of GATA1. Binds to hormone response elements (HREs) consisting of two 5'-AGGTCA-3' half site direct repeat consensus sequences. Plays a fundamental role in early embryonic development and embryonic stem cells. Required for normal spermatogenesis and cerebellum development. Appears to be important for neurodevelopmentally regulated behavior (By similarity). Activates transcriptional activity of LHCG. Antagonist of PPARA-mediated transactivation.

**Cellular Location**

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00407, ECO:0000269|PubMed:10644740, ECO:0000269|PubMed:15302918}

## NR2C2/TR4 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## NR2C2/TR4 Rabbit mAb - Images

