

Phospho-ULK1 (Ser556) Rabbit mAb
Catalog # AP78924**Specification****Phospho-ULK1 (Ser556) Rabbit mAb - Product Information**

| | |
|-------------------|------------------------|
| Application | WB |
| Primary Accession | O75385 |
| Reactivity | Human, Mouse |
| Host | Rabbit |
| Clonality | Monoclonal Antibody |
| Calculated MW | 112631 |

Phospho-ULK1 (Ser556) Rabbit mAb - Additional Information

Gene ID 8408

Other Names

ULK1

Dilution

WB~~1/500-1/1000

Format

Liquid

Phospho-ULK1 (Ser556) Rabbit mAb - Protein Information**Name** ULK1 {ECO:0000303|PubMed:9693035, ECO:0000312|HGNC:HGNC:12558}**Function**

Serine/threonine-protein kinase involved in autophagy in response to starvation (PubMed: 18936157, PubMed: 21460634, PubMed: 21795849, PubMed: 23524951, PubMed: 25040165, PubMed: 29487085, PubMed: 31123703). Acts upstream of phosphatidylinositol 3-kinase PIK3C3 to regulate the formation of autophagophores, the precursors of autophagosomes (PubMed: 18936157, PubMed: 21460634, PubMed: 21795849, PubMed: 25040165). Part of regulatory feedback loops in autophagy: acts both as a downstream effector and negative regulator of mammalian target of rapamycin complex 1 (mTORC1) via interaction with RPTOR (PubMed: 21795849). Activated via phosphorylation by AMPK and also acts as a regulator of AMPK by mediating phosphorylation of

AMPK subunits PRKAA1, PRKAB2 and PRKAG1, leading to negatively regulate AMPK activity (PubMed:21460634). May phosphorylate ATG13/KIAA0652 and RPTOR; however such data need additional evidences (PubMed:18936157). Plays a role early in neuronal differentiation and is required for granule cell axon formation (PubMed:11146101). Also phosphorylates SESN2 and SQSTM1 to regulate autophagy (PubMed:25040165, PubMed:37306101). Phosphorylates FLCN, promoting autophagy (PubMed:25126726). Phosphorylates AMBRA1 in response to autophagy induction, releasing AMBRA1 from the cytoskeletal docking site to induce autophagosome nucleation (PubMed:20921139). Phosphorylates ATG4B, leading to inhibit autophagy by decreasing both proteolytic activation and delipidation activities of ATG4B (PubMed:28821708).

Cellular Location

Cytoplasm, cytosol. Preautophagosomal structure. Note=Under starvation conditions, is localized to punctate structures primarily representing the isolation membrane that sequesters a portion of the cytoplasm resulting in the formation of an autophagosome.

Tissue Location

Ubiquitously expressed. Detected in the following adult tissues: skeletal muscle, heart, pancreas, brain, placenta, liver, kidney, and lung

Phospho-ULK1 (Ser556) Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Phospho-ULK1 (Ser556) Rabbit mAb - Images



