

**PKM2 Rabbit mAb**  
Catalog # AP78976**Specification**

---

**PKM2 Rabbit mAb - Product Information**

Application	<b>WB</b>
Primary Accession	<a href="#">P14618</a>
Reactivity	<b>Human, Mouse, Rat</b>
Host	<b>Rabbit</b>
Clonality	<b>Monoclonal Antibody</b>
Calculated MW	<b>57937</b>

**PKM2 Rabbit mAb - Additional Information****Gene ID** 5315**Other Names**

PKM

**Dilution**

WB~~1/500-1/1000

**Format**

Liquid

**PKM2 Rabbit mAb - Protein Information****Name** PKM**Synonyms** OIP3 {ECO:0000303|PubMed:9466265}, PK2,**Function**

Catalyzes the final rate-limiting step of glycolysis by mediating the transfer of a phosphoryl group from phosphoenolpyruvate (PEP) to ADP, generating ATP (PubMed: [15996096](http://www.uniprot.org/citations/15996096), PubMed: [1854723](http://www.uniprot.org/citations/1854723), PubMed: [20847263](http://www.uniprot.org/citations/20847263)). The ratio between the highly active tetrameric form and nearly inactive dimeric form determines whether glucose carbons are channeled to biosynthetic processes or used for glycolytic ATP production (PubMed: [15996096](http://www.uniprot.org/citations/15996096), PubMed: [1854723](http://www.uniprot.org/citations/1854723), PubMed: [20847263](http://www.uniprot.org/citations/20847263)). The transition between the 2 forms contributes to the control of glycolysis and is important for tumor cell proliferation and survival (PubMed: [15996096](http://www.uniprot.org/citations/15996096), PubMed: [1854723](http://www.uniprot.org/citations/1854723), PubMed: [20847263](http://www.uniprot.org/citations/20847263)).

### Cellular Location

[Isoform M2]: Cytoplasm. Nucleus Note=Translocates to the nucleus in response to various signals, such as EGF receptor activation or apoptotic stimuli (PubMed:17308100, PubMed:22056988, PubMed:24120661). Nuclear translocation is promoted by acetylation by EP300 (PubMed:24120661). Deacetylation by SIRT6 promotes its nuclear export in a process dependent of XPO4, thereby suppressing its ability to activate transcription and promote tumorigenesis (PubMed:26787900).

### Tissue Location

[Isoform M2]: Specifically expressed in proliferating cells, such as embryonic stem cells, embryonic carcinoma cells, as well as cancer cells.

### PKM2 Rabbit mAb - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### PKM2 Rabbit mAb - Images

