

**CYP7A1 Antibody (C-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP7996A****Specification**

---

**CYP7A1 Antibody (C-term) - Product Information**

Application	<b>WB,E</b>
Primary Accession	<a href="#">P22680</a>
Other Accession	<a href="#">P51542</a>
Reactivity	<b>Human</b>
Predicted	<b>Rabbit</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Isotype	<b>Rabbit IgG</b>
Calculated MW	<b>57661</b>
Antigen Region	<b>476-504</b>

**CYP7A1 Antibody (C-term) - Additional Information****Gene ID** 1581**Other Names**

Cholesterol 7-alpha-monooxygenase, CYPVII, Cholesterol 7-alpha-hydroxylase, Cytochrome P450 7A1, CYP7A1, CYP7

**Target/Specificity**

This CYP7A1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 476-504 amino acids from the C-terminal region of human CYP7A1.

**Dilution**

WB~~1:1000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

CYP7A1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**CYP7A1 Antibody (C-term) - Protein Information****Name** CYP7A1 {ECO:0000303|PubMed:12077124, ECO:0000312|HGNC:HGNC:2651}

**Function** A cytochrome P450 monooxygenase involved in the metabolism of endogenous cholesterol and its oxygenated derivatives (oxysterols) (PubMed:[11013305](#), PubMed:[12077124](#), PubMed:[19965590](#), PubMed:[21813643](#), PubMed:[2384150](#)). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (CPR; NADPH-ferrihemoprotein reductase) (PubMed:[11013305](#), PubMed:[12077124](#), PubMed:[19965590](#), PubMed:[21813643](#), PubMed:[2384150](#)). Functions as a critical regulatory enzyme of bile acid biosynthesis and cholesterol homeostasis. Catalyzes the hydroxylation of carbon hydrogen bond at 7-alpha position of cholesterol, a rate-limiting step in cholesterol catabolism and bile acid biosynthesis (PubMed:[12077124](#), PubMed:[19965590](#), PubMed:[2384150](#)). 7-alpha hydroxylates several oxysterols, including 4beta-hydroxycholesterol and 24-hydroxycholesterol (PubMed:[11013305](#), PubMed:[12077124](#)). Catalyzes the oxidation of the 7,8 double bond of 7-dehydrocholesterol and lathosterol with direct and predominant formation of the 7-keto derivatives (PubMed:[21813643](#)).

#### Cellular Location

Endoplasmic reticulum membrane; Single-pass membrane protein. Microsome membrane; Single-pass membrane protein

#### Tissue Location


Detected in liver..

### CYP7A1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### CYP7A1 Antibody (C-term) - Images



Western blot analysis of CYP7A1 Antibody (C-term) (Cat.#AP7996a) in K562(lane 1), HepG2(lane

2) cell line lysates (35ug/lane). CYP7A1 (arrow) was detected using the purified Pab.



Anti-CYP7A1 Antibody (C-term) at 1:1000 dilution + HepG2 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 58 kDa Blocking/Dilution buffer: 5% NFDm/TBST.

#### **CYP7A1 Antibody (C-term) - Background**

CYP7A1 is a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This endoplasmic reticulum membrane protein catalyzes the first reaction in the cholesterol catabolic pathway in the liver, which converts cholesterol to bile acids. This reaction is the rate limiting step and the major site of regulation of bile acid synthesis, which is the primary mechanism for the removal of cholesterol from the body.

#### **CYP7A1 Antibody (C-term) - References**

Lenicek, M., J. Lipid Res. 49 (12), 2664-2667 (2008)  
Nelson, D.R., Pharmacogenetics 14 (1), 1-18 (2004)