

BRD4 Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP8051a

Specification

BRD4 Antibody (N-term) - Product Information

Application	IHC-P,E
Primary Accession	O60885
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	1-30

BRD4 Antibody (N-term) - Additional Information

Gene ID 23476

Other Names

Bromodomain-containing protein 4, Protein HUNK1, BRD4, HUNK1

Target/Specificity

This BRD4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1~30 amino acids from the N-terminal region of human BRD4.

Dilution

IHC-P~~1:50~100

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

BRD4 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

BRD4 Antibody (N-term) - Protein Information

Name BRD4

Synonyms HUNK1

Function Chromatin reader protein that recognizes and binds acetylated histones and plays a key role in transmission of epigenetic memory across cell divisions and transcription regulation

(PubMed:[20871596](#), PubMed:[23086925](#), PubMed:[23317504](#), PubMed:[29176719](#)). Remains associated with acetylated chromatin throughout the entire cell cycle and provides epigenetic memory for postmitotic G1 gene transcription by preserving acetylated chromatin status and maintaining high-order chromatin structure (PubMed:[22334664](#), PubMed:[23317504](#), PubMed:[23589332](#)). During interphase, plays a key role in regulating the transcription of signal-inducible genes by associating with the P-TEFb complex and recruiting it to promoters (PubMed:[16109376](#), PubMed:[16109377](#), PubMed:[19596240](#), PubMed:[23589332](#), PubMed:[24360279](#)). Also recruits P-TEFb complex to distal enhancers, so called anti-pause enhancers in collaboration with JMJD6 (PubMed:[16109376](#), PubMed:[16109377](#), PubMed:[19596240](#), PubMed:[23589332](#), PubMed:[24360279](#)). BRD4 and JMJD6 are required to form the transcriptionally active P-TEFb complex by displacing negative regulators such as HEXIM1 and 7SKsnRNA complex from P-TEFb, thereby transforming it into an active form that can then phosphorylate the C-terminal domain (CTD) of RNA polymerase II (PubMed:[16109376](#), PubMed:[16109377](#), PubMed:[19596240](#), PubMed:[23589332](#), PubMed:[24360279](#)). Regulates differentiation of naive CD4(+) T-cells into T-helper Th17 by promoting recruitment of P-TEFb to promoters (By similarity). Promotes phosphorylation of 'Ser-2' of the C-terminal domain (CTD) of RNA polymerase II (PubMed:[23086925](#)). According to a report, directly acts as an atypical protein kinase and mediates phosphorylation of 'Ser-2' of the C-terminal domain (CTD) of RNA polymerase II; these data however need additional evidences in vivo (PubMed:[22509028](#)). In addition to acetylated histones, also recognizes and binds acetylated RELA, leading to further recruitment of the P-TEFb complex and subsequent activation of NF-kappa-B (PubMed:[19103749](#)). Also acts as a regulator of p53/TP53-mediated transcription: following phosphorylation by CK2, recruited to p53/TP53 specific target promoters (PubMed:[23317504](#)).

Cellular Location

Nucleus. Chromosome. Note=Associates with acetylated chromatin (PubMed:[16109376](#), PubMed:[21890894](#)). Released from chromatin upon deacetylation of histones that can be triggered by different signals such as activation of the JNK pathway or nocodazole treatment (PubMed:[16109376](#), PubMed:[21890894](#)). Preferentially localizes to mitotic chromosomes, while it does not localize to meiotic chromosomes (PubMed:[16109376](#), PubMed:[21890894](#)).

Tissue Location

Ubiquitously expressed.


BRD4 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BRD4 Antibody (N-term) - Images





Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by AEC staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

BRD4 Antibody (N-term) - Background

BRD4 is homologous to the murine protein MCAP, which associates with chromosomes during mitosis, and to the human RING3 protein, a serine/threonine kinase. Each of these proteins contains two bromodomains, a conserved sequence motif which may be involved in chromatin targeting. The gene has been implicated as the chromosome 19 target of translocation t(15;19)(q13;p13.1), which defines an upper respiratory tract carcinoma in young people.

BRD4 Antibody (N-term) - References

Gagnon, D. et al. J Virol. May; 83(9): 4127-4139(2009).
Maruyama, T., et al., Mol. Cell. Biol. 22(18):6509-6520 (2002).
French, C.A., et al., Am. J. Pathol. 159(6):1987-1992 (2001).
Dey, A., et al., Mol. Cell. Biol. 20(17):6537-6549 (2000).

BRD4 Antibody (N-term) - Citations

- [Proteasomal degradation of the papillomavirus E2 protein is inhibited by overexpression of bromodomain-containing protein 4.](#)