

**TRPM7 (CHAK1) Antibody (N-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP8052a**

**Specification**

---

**TRPM7 (CHAK1) Antibody (N-term) - Product Information**

Application	IHC-P,E
Primary Accession	<a href="#">O96QT4</a>
Other Accession	<a href="#">O9BXB2</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	45-74

**TRPM7 (CHAK1) Antibody (N-term) - Additional Information**

**Gene ID** 54822

**Other Names**

Transient receptor potential cation channel subfamily M member 7, Channel-kinase 1, Long transient receptor potential channel 7, LTrpC-7, LTrpC7, TRPM7, CHAK1, LTRPC7

**Target/Specificity**

This TRPM7 (CHAK1) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 45-74 amino acids from the N-terminal region of human TRPM7 (CHAK1).

**Dilution**

IHC-P~~1:50~100

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

TRPM7 (CHAK1) Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**TRPM7 (CHAK1) Antibody (N-term) - Protein Information**

**Name** TRPM7

**Synonyms** CHAK1, LTRPC7 {ECO:0000303|PubMed:113855

**Function** Bifunctional protein that combines an ion channel with an intrinsic kinase domain, enabling it to modulate cellular functions either by conducting ions through the pore or by phosphorylating downstream proteins via its kinase domain. The channel is highly permeable to divalent cations, specifically calcium (Ca<sup>2+</sup>), magnesium (Mg<sup>2+</sup>) and zinc (Zn<sup>2+</sup>) and mediates their influx (PubMed:[11385574](#), PubMed:[12887921](#), PubMed:[15485879](#), PubMed:[24316671](#), PubMed:[35561741](#), PubMed:[36027648](#)). Controls a wide range of biological processes such as Ca<sup>2+</sup>(+), Mg<sup>2+</sup>(+) and Zn<sup>2+</sup>(+) homeostasis, vesicular Zn<sup>2+</sup>(+) release channel and intracellular Ca<sup>2+</sup>(+) signaling, embryonic development, immune responses, cell motility, proliferation and differentiation (By similarity). The C-terminal alpha-kinase domain autophosphorylates cytoplasmic residues of TRPM7 (PubMed:[18365021](#)). In vivo, TRPM7 phosphorylates SMAD2, suggesting that TRPM7 kinase may play a role in activating SMAD signaling pathways. In vitro, TRPM7 kinase phosphorylates ANXA1 (annexin A1), myosin II isoforms and a variety of proteins with diverse cellular functions (PubMed:[15485879](#), PubMed:[18394644](#)).

#### Cellular Location


Cell membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:Q923J1}. Cytoplasmic vesicle membrane {ECO:0000250|UniProtKB:Q923J1}; Multi-pass membrane protein {ECO:0000250|UniProtKB:Q923J1}. Note=Localized largely in intracellular Zn<sup>2+</sup>(+)-storage vesicles. {ECO:0000250|UniProtKB:Q923J1}

#### TRPM7 (CHAK1) Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### TRPM7 (CHAK1) Antibody (N-term) - Images



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

### **TRPM7 (CHAK1) Antibody (N-term) - Background**

TRPCs, mammalian homologs of the *Drosophila* transient receptor potential (*trp*) protein, are ion channels that are thought to mediate capacitative calcium entry into the cell. TRP-PLIK is a protein that is both an ion channel and a kinase. As a channel, it conducts calcium and monovalent cations to depolarize cells and increase intracellular calcium. As a kinase, it is capable of phosphorylating itself and other substrates. The kinase activity is necessary for channel function, as shown by its dependence on intracellular ATP and by the kinase mutants.[supplied by OMIM]

### **TRPM7 (CHAK1) Antibody (N-term) - References**

Blume-Jensen P, et al. *Nature* 2001. 411: 355.  
Cantrell D, J. *Cell Sci.* 2001. 114: 1439.  
Jiang S *Oncogene* 2000. 19: 5590.  
Manning G, et al. *Science* 2002. 298: 1912.  
Moller, D, et al. *Am. J. Physiol.* 1994. 266: C351-C359.  
Robertson, S. et al. *Trends Genet.* 2000. 16: 368.  
Robinson D, et al. *Oncogene* 2000. 19: 5548.  
Van der Ven, P, et al. *Hum. Molec. Genet.* 1993. 2: 1889.  
Vanhaesebroeck, B, et al. *Biochem. J.* 2000. 346: 561.  
Van Weering D, et al. *Recent Results Cancer Res.* 1998. 154: 271.

### **TRPM7 (CHAK1) Antibody (N-term) - Citations**

- [Transient receptor potential melastatin 7-like current in human head and neck carcinoma cells: role in cell proliferation.](#)