

NEK3 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP8075c

Specification

NEK3 Antibody (Center) - Product Information

| | |
|-------------------|---------------------------|
| Application | WB,E |
| Primary Accession | P51956 |
| Other Accession | NP_002489 |
| Reactivity | Human |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | Rabbit IgG |
| Antigen Region | 331-361 |

NEK3 Antibody (Center) - Additional Information

Gene ID 4752

Other Names

Serine/threonine-protein kinase Nek3, HSPK 36, Never in mitosis A-related kinase 3, NimA-related protein kinase 3, NEK3

Target/Specificity

This NEK3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 331-361 amino acids from the Central region of human NEK3.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

NEK3 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

NEK3 Antibody (Center) - Protein Information

Name NEK3

Function Protein kinase which influences neuronal morphogenesis and polarity through effects on microtubules. Regulates microtubule acetylation in neurons. Contributes to prolactin-mediated

phosphorylation of PXN and VAV2. Implicated in prolactin-mediated cytoskeletal reorganization and motility of breast cancer cells through mechanisms involving RAC1 activation and phosphorylation of PXN and VAV2.

Cellular Location

Cytoplasm. Cell projection, axon

Tissue Location


Up-regulated in malignant versus normal breast tissue. Isoform 2 shows a high level of expression in testis, ovary and brain.

NEK3 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


NEK3 Antibody (Center) - Images



All lanes : Anti-NEK3 Antibody (K346) at 1:2000 dilution Lane 1: A431 whole cell lysate Lane 2: MCF-7 whole cell lysate Lane 3: SH-SY5Y whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 58 kDa Blocking/Dilution buffer: 5% NFDN/TBST.



All lanes : Anti-NEK3 Antibody (K346) at 1:2000 dilution Lane 1: HepG2 whole cell lysate Lane 2: CCRF-CEM whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 58 kDa Blocking/Dilution buffer: 5% NFDN/TBST.



Western blot analysis of NEK3 (arrow) using rabbit polyclonal NEK3 Antibody (K346) (Cat. #AP8075c). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the NEK3 gene.

NEK3 Antibody (Center) - Background

In *Aspergillus nidulans*, lack of the serine/threonine kinase NimA (never in mitosis A) results in cell cycle arrest in G2, while overexpression causes the premature onset of mitotic events. NEK3 is similar in sequence to the *Aspergillus nidulans* protein and may therefore play a role in mitotic regulation. However, the encoded protein differs from other NimA family members in that it is not cell cycle regulated and is found primarily in the cytoplasm.

NEK3 Antibody (Center) - References

Schultz, S.J., et al., *Cell Growth Differ.* 5(6):625-635 (1994).
Schultz, S.J., et al., *Cell Growth Differ.* 4(10):821-830 (1993).
Kimura, M., et al., *Cytogenet. Cell Genet.* 95 (3-4), 177-182 (2001).

NEK3 Antibody (Center) - Citations

- [Overexpression of NEK3 is associated with poor prognosis in patients with gastric cancer.](#)
- [Altered expression of prolactin receptor-associated signaling proteins in human breast](#)

[carcinoma.](#)