

**PACSIN2 Antibody (C-term)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP8088b**

**Specification**

---

**PACSIN2 Antibody (C-term) - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | <b>WB, IHC-P,E</b>     |
| Primary Accession | <a href="#">O9UNF0</a> |
| Reactivity        | <b>Human, Mouse</b>    |
| Host              | <b>Rabbit</b>          |
| Clonality         | <b>Polyclonal</b>      |
| Isotype           | <b>Rabbit IgG</b>      |
| Antigen Region    | <b>342-371</b>         |

**PACSIN2 Antibody (C-term) - Additional Information**

**Gene ID** 11252

**Other Names**

Protein kinase C and casein kinase substrate in neurons protein 2, Syndapin-2, Syndapin-II, PACSIN2

**Target/Specificity**

This PACSIN2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 342-371 amino acids from the C-terminal region of human PACSIN2.

**Dilution**

WB~~1:1000  
IHC-P~~1:50~100

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

PACSIN2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**PACSIN2 Antibody (C-term) - Protein Information**

**Name** PACSIN2

**Function** Regulates the morphogenesis and endocytosis of caveolae (By similarity). Lipid-binding protein that is able to promote the tubulation of the phosphatidic acid-containing membranes it

preferentially binds. Plays a role in intracellular vesicle-mediated transport. Involved in the endocytosis of cell-surface receptors like the EGF receptor, contributing to its internalization in the absence of EGF stimulus.

#### Cellular Location

Cytoplasm. Cytoplasm, cytoskeleton. Cytoplasmic vesicle membrane; Peripheral membrane protein; Cytoplasmic side. Early endosome. Recycling endosome membrane Cell projection, ruffle membrane; Peripheral membrane protein; Cytoplasmic side. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection Membrane, caveola. Note=Detected at the neck of flask-shaped caveolae. Localization to tubular recycling endosomes probably requires interaction with MICALL1 and EHD1

#### Tissue Location


Widely expressed.

### PACSIN2 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### PACSIN2 Antibody (C-term) - Images



PACSIN2 Antibody (S357) (Cat.# AP8088b) western blot analysis in Daudi,HL-60 cell line and mouse brain,rat heart lysates (35ug/lane).This demonstrates the PACSIN2 antibody detected the PACSIN2 protein (arrow).



Immunohistochemical analysis of paraffin-embedded H. stomach section using PACSIN2 Antibody (C-term)(Cat#AP8088b). AP8088b was diluted at 1:25 dilution. A peroxidase-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody, followed by DAB staining.



Formalin-fixed and paraffin-embedded human cancer tissue reacted with the primary antibody, which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated. BC = breast carcinoma; HC = hepatocarcinoma.

### **PACSIN2 Antibody (C-term) - Background**

PACSIN may play a role in vesicle formation and transport. This protein homo- and hetero-aggregates with other PACSINs. It also binds dynamin 1, synaptojanin, synapsin 1 and the neural Wiskott-Aldrich syndrome protein (N-WASP). The protein exhibits a vesicle-like cytoplasmic distribution and is ubiquitously expressed. PACSIN is phosphorylated by casein kinase 2 (CK2) and protein kinase C (PKC). The protein contains 1 FCH domain and 1 SH3 domain.

### **PACSIN2 Antibody (C-term) - References**

Strausberg, R.L., et al., Proc. Natl. Acad. Sci. U.S.A. 99(26):16899-16903 (2002). Wiemann, S., et al., Genome Res. 11(3):422-435 (2001). Ritter, B., et al., FEBS Lett. 454(3):356-362 (1999). Dunham, I., et al., Nature 402(6761):489-495 (1999).

**PACSIN2 Antibody (C-term) - Citations**

- [Expression of a Human Caveolin-1 Mutation in Mice Drives Inflammatory and Metabolic Defect-Associated Pulmonary Arterial Hypertension.](#)
- [A disease-associated frameshift mutation in caveolin-1 disrupts caveolae formation and function through introduction of a de novo ER retention signal.](#)
- [Clostridium difficile Toxin A Undergoes Clathrin-Independent, PACSIN2-Dependent Endocytosis.](#)
- [Characterization of a caveolin-1 mutation associated with both PAH and congenital generalized lipodystrophy.](#)
- [EHD3 is Required for Tubular Recycling Endosome Stabilization and an Asparagine-Glutamic Acid Residue Pair within its EH Domain Dictates its Selective Binding to NPF Peptides.](#)
- [Endocytic recycling protein EHD1 regulates primary cilia morphogenesis and SHH signaling during neural tube development.](#)
- [Differential roles of C-terminal Eps15 homology domain proteins as vesiculators and tubulators of recycling endosomes.](#)
- [Cooperation of MICAL-L1, syndapin2, and phosphatidic acid in tubular recycling endosome biogenesis.](#)
- [The F-BAR protein PACSIN2 regulates epidermal growth factor receptor internalization.](#)
- [Pacsin 2 is recruited to caveolae and functions in caveolar biogenesis.](#)
- [The F-BAR domain protein PACSIN2 associates with Rac1 and regulates cell spreading and migration.](#)