

### **ADH5 Antibody (Center)**

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP8562C

### Specification

# ADH5 Antibody (Center) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	<u>P11766</u>
Other Accession	<u>019053</u>
Reactivity	Human
Predicted	Rabbit
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	212-239

## ADH5 Antibody (Center) - Additional Information

Gene ID 128

**Other Names** 

Alcohol dehydrogenase class-3, Alcohol dehydrogenase 5, Alcohol dehydrogenase class chi chain, Alcohol dehydrogenase class-III, Glutathione-dependent formaldehyde dehydrogenase, FALDH, FDH, GSH-FDH, 111-, S-(hydroxymethyl)glutathione dehydrogenase, ADH5 (<a href="http://www.genenames.org/cgi-bin/gene\_symbol\_report?hgnc\_id=253" target="\_blank">HGNC:253</a>), ADHX, FDH

### Target/Specificity

This ADH5 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 212-239 amino acids from the Central region of human ADH5.

**Dilution** WB~~1:4000 IHC-P~~1:100 FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### Precautions

ADH5 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

### **ADH5 Antibody (Center) - Protein Information**



Name ADH5 (<u>HGNC:253</u>)

Synonyms ADHX, FDH

**Function** Catalyzes the oxidation of long-chain primary alcohols and the oxidation of S-(hydroxymethyl) glutathione (PubMed:<u>8460164</u>). Also oxidizes long chain omega-hydroxy fatty acids, such as 20-HETE, producing both the intermediate aldehyde, 20-oxoarachidonate and the end product, a dicarboxylic acid, (5Z,8Z,11Z,14Z)-eicosatetraenedioate (PubMed:<u>16081420</u>). Class-III ADH is remarkably ineffective in oxidizing ethanol (PubMed:<u>8460164</u>). Required for clearance of cellular formaldehyde, a cytotoxic and carcinogenic metabolite that induces DNA damage (PubMed:<u>33355142</u>). Also acts as a S-nitroso-glutathione reductase by catalyzing the NADH-dependent reduction of S- nitrosoglutathione, thereby regulating protein S-nitrosylation (By similarity).

Cellular Location Cytoplasm.

## ADH5 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

### ADH5 Antibody (Center) - Images



Western blot analysis of ADH5 (arrow) using rabbit polyclonal ADH5 Antibody (Center) (Cat. #AP8562c). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the ADH5 gene.





All lanes : Anti-ADH5 Antibody (Center) at 1:4000 dilution Lane 1: K562 whole cell lysate Lane 2: SH-SY5Y whole cell lysate Lysates/proteins at 20  $\mu$ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 40 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Formalin-fixed and paraffin-embedded human hepatocarcinoma with ADH5 Antibody (Center), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



Formalin-fixed and paraffin-embedded H.breast tissue reacted with ADH5 Antibody (Center)



## (Cat#AP8562c).



Formalin-fixed and paraffin-embedded H.placenta tissue reacted with ADH5 Antibody (Center) (Cat#AP8562c).



Flow cytometric analysis of K562 cells using ADH5 Antibody (Center)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

## ADH5 Antibody (Center) - Background

ADH5 is a member of the alcohol dehydrogenase family. Members of this family metabolize a wide variety of substrates, including ethanol, retinol, other aliphatic alcohols, hydroxysteroids, and lipid peroxidation products. This protein forms a homodimer. It has virtually no activity for ethanol oxidation, but exhibits high activity for oxidation of long-chain primary alcohols and for oxidation of S-hydroxymethyl-glutathione, a spontaneous adduct between formaldehyde and glutathione. This enzyme is an important component of cellular metabolism for the elimination of formaldehyde, a potent irritant and sensitizing agent that causes lacrymation, rhinitis, pharyngitis, and contact dermatitis.

## **ADH5 Antibody (Center) - References**

Martins-de-Souza, D., et.al., Eur Arch Psychiatry Clin Neurosci 259 (3), 151-163 (2009) Iborra, F.J., et.al., J. Histochem. Cytochem. 40 (12), 1865-1878 (1992)