

IGFBP2 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP8588B**Specification**

IGFBP2 Antibody (C-term) - Product Information

Application	IF, WB, IHC-P, FC,E
Primary Accession	P18065
Other Accession	P24853 , P13384 , Q29400
Reactivity	Human
Predicted	Bovine, Pig, Sheep
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	277-305

IGFBP2 Antibody (C-term) - Additional Information**Gene ID** 3485**Other Names**

Insulin-like growth factor-binding protein 2, IBP-2, IGF-binding protein 2, IGFBP-2, IGFBP2, BP2, IBP2

Target/Specificity

This IGFBP2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 277-305 amino acids from the C-terminal region of human IGFBP2.

DilutionIF~~1:10~50
WB~~1:1000
IHC-P~~1:10~50
FC~~1:10~50**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

IGFBP2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

IGFBP2 Antibody (C-term) - Protein Information

Name IGFBP2

Synonyms BP2, IBP2

Function Inhibits IGF-mediated growth and developmental rates. IGF- binding proteins prolong the half-life of the IGFs and have been shown to either inhibit or stimulate the growth promoting effects of the IGFs on cell culture. They alter the interaction of IGFs with their cell surface receptors.


Cellular Location
Secreted.

IGFBP2 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


IGFBP2 Antibody (C-term) - Images




Confocal immunofluorescent analysis of IGF2BP2 Antibody (C-term)(Cat. #AP8588b) with A549 cell followed by Alexa Fluor[®]488-conjugated goat anti-rabbit IgG (green). DAPI was used to stain the cell nuclear (blue).



Anti-IGFBP2 Antibody (C-term) at 1:1000 dilution + T47D whole cell lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 35 kDa Blocking/Dilution buffer: 5% NFDN/TBST.



Formalin-fixed and paraffin-embedded human brain tissue with IGFBP2 Antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



Flow cytometric analysis of jurkat cells using IGFBP2 Antibody (C-term)(bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

IGFBP2 Antibody (C-term) - Background

IGFBP2-binding proteins prolong the half-life of the IGFs and have been shown to either inhibit or stimulate the growth promoting effects of the IGFs on cell culture. They alter the interaction of IGFs with their cell surface receptors.

IGFBP2 Antibody (C-term) - References

- Arafat,A.M.,et.al.,J. Clin. Endocrinol. Metab. 94 (12), 5093-5101 (2009)
- Yazawa,T.,et.al.,Am. J. Pathol. 175 (3), 976-987 (2009)