

CYP2E1 Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP8644B

Specification

CYP2E1 Antibody (C-term) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	P05181
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	56849
Antigen Region	402-429

CYP2E1 Antibody (C-term) - Additional Information

Gene ID 1571

Other Names

Cytochrome P450 2E1, 11413-, 4-nitrophenol 2-hydroxylase, 11413n7, CYP11E1, Cytochrome P450-J, Cytochrome P450 2E1, N-terminally processed, CYP2E1, CYP2E

Target/Specificity

This CYP2E1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 402-429 amino acids from the C-terminal region of human CYP2E1.

Dilution

WB~~1:1000
IHC-P~~1:50~100
FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CYP2E1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

CYP2E1 Antibody (C-term) - Protein Information

Name CYP2E1 {ECO:0000303|PubMed:10553002, ECO:0000312|HGNC:HGNC:2631}

Function A cytochrome P450 monooxygenase involved in the metabolism of fatty acids (PubMed:[10553002](#), PubMed:[18577768](#)). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate, and reducing the second into a water molecule, with two electrons provided by NADPH via cytochrome P450 reductase (NADPH--hemoprotein reductase) (PubMed:[10553002](#), PubMed:[18577768](#)). Catalyzes the hydroxylation of carbon-hydrogen bonds. Hydroxylates fatty acids specifically at the omega-1 position displaying the highest catalytic activity for saturated fatty acids (PubMed:[10553002](#), PubMed:[18577768](#)). May be involved in the oxidative metabolism of xenobiotics (Probable).

Cellular Location


Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:P05182}; Peripheral membrane protein {ECO:0000250|UniProtKB:P05182}. Microsome membrane {ECO:0000250|UniProtKB:P05182}; Peripheral membrane protein {ECO:0000250|UniProtKB:P05182}. Mitochondrion inner membrane {ECO:0000250|UniProtKB:P05182}; Peripheral membrane protein {ECO:0000250|UniProtKB:P05182}. Note=Post-translationally targeted to mitochondria. TOMM70 is required for the translocation across the mitochondrial outer membrane. After translocation into the matrix, associates with the inner membrane as a membrane extrinsic protein {ECO:0000250|UniProtKB:P05182}

CYP2E1 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


CYP2E1 Antibody (C-term) - Images



Western blot analysis of CYP2E1 Antibody (C-term) (Cat. #AP8644b) in K562 cell line lysates (35ug/lane). CYP2E1 (arrow) was detected using the purified Pab.



Formalin-fixed and paraffin-embedded human brain tissue reacted with CYP2E1 Antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



CYP2E1 Antibody (C-term) (Cat.#AP8644b) flow cytometry analysis of HepG2 cells (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

CYP2E1 Antibody (C-term) - Background

CYP2E1 is a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and is induced by ethanol, the diabetic state, and starvation. The enzyme metabolizes both endogenous substrates, such as ethanol, acetone, and acetal, as well as exogenous substrates including benzene, carbon tetrachloride, ethylene glycol, and nitrosamines which are premutagens found in cigarette smoke. Due to its many substrates, this enzyme may be involved in such varied processes as gluconeogenesis, hepatic cirrhosis, diabetes, and cancer.

CYP2E1 Antibody (C-term) - References

Mingazova,S.R., et.al., Med Tr Prom Ekol 11, 30-33 (2009)
Robinson,R.C., et.al., Pharmacology 39 (3), 137-144 (1989)