

HSPA7 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP8748C**Specification**

HSPA7 Antibody (Center) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	P48741
Other Accession	P17066
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	169-196

HSPA7 Antibody (Center) - Additional Information**Other Names**

Putative heat shock 70 kDa protein 7, Heat shock 70 kDa protein B, HSPA7, HSP70B

Target/Specificity

This HSPA7 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 169-196 amino acids from the Central region of human HSPA7.

Dilution

WB~~1:2000
IHC-P~~1:250
FC~~1:25

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

HSPA7 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

HSPA7 Antibody (Center) - Protein Information

Name HSPA7


Synonyms HSP70B

HSPA7 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HSPA7 Antibody (Center) - Images




Anti-HSPA7 Antibody (Center) at 1:2000 dilution + 293 whole cell lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 70 kDa Blocking/Dilution buffer: 5% NFDN/TBST.




Anti-HSPA7 Antibody (Center) at 1:2000 dilution + CCRF-CEM whole cell lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000


dilution. Predicted band size : 70 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



AP8748C staining HSPA7 in human lung adenocarcinoma tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Samples were incubated with primary antibody (1/250) for 1 hours at room temperature. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.



AP8748C staining HSPA7 in human testis tissue sections by Immunohistochemistry (IHC-P - paraformaldehyde-fixed, paraffin-embedded sections). Samples were incubated with primary antibody (1/250) for 1 hours at room temperature. A undiluted biotinylated goat polyvalent antibody was used as the secondary antibody.



Overlay histogram showing U-2 OS cells stained with AP8748C(green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AP8748C, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed(1583138) at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was rabbit IgG1 (1µg/1x10⁶ cells) used under the same conditions. Acquisition of >10, 000 events was performed.

HSPA7 Antibody (Center) - References

Voellmy R., et.al., Proc. Natl. Acad. Sci. U.S.A. 82:4949-4953(1985).