

**CLIP1 Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP8950a**

**Specification**

---

**CLIP1 Antibody (N-term) - Product Information**

Application	WB, IHC-P, FC,E
Primary Accession	<a href="#">P30622</a>
Other Accession	<a href="#">O55156</a> , <a href="#">O9Z0H8</a> , <a href="#">O9UDT6</a> , <a href="#">O922J3</a> , <a href="#">O42184</a> , <a href="#">O9JK25</a>
Reactivity	Human
Predicted	Chicken, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	162246
Antigen Region	228-254

**CLIP1 Antibody (N-term) - Additional Information**

**Gene ID** 6249

**Other Names**

CAP-Gly domain-containing linker protein 1, Cytoplasmic linker protein 1, Cytoplasmic linker protein 170 alpha-2, CLIP-170, Reed-Sternberg intermediate filament-associated protein, Restin, CLIP1, CYLN1, RSN

**Target/Specificity**

This CLIP1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 228-254 amino acids from the N-terminal region of human CLIP1.

**Dilution**

WB~~1:1000  
IHC-P~~1:50~100  
FC~~1:10~50

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

CLIP1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**CLIP1 Antibody (N-term) - Protein Information**

## Name CLIP1

## Synonyms CYLN1, RSN

**Function** Binds to the plus end of microtubules and regulates the dynamics of the microtubule cytoskeleton. Promotes microtubule growth and microtubule bundling. Links cytoplasmic vesicles to microtubules and thereby plays an important role in intracellular vesicle trafficking. Plays a role in macropinocytosis and endosome trafficking.

## Cellular Location

Cytoplasm. Cytoplasm, cytoskeleton. Cytoplasmic vesicle membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection, ruffle. Note=Localizes to microtubule plus ends (PubMed:17889670, PubMed:21646404). Localizes preferentially to the ends of tyrosinated microtubules (By similarity). Accumulates in plasma membrane regions with ruffling and protrusions. Associates with the membranes of intermediate macropinocytic vesicles (PubMed:12433698) {ECO:0000250|UniProtKB:Q922J3, ECO:0000269|PubMed:12433698, ECO:0000269|PubMed:17889670, ECO:0000269|PubMed:21646404}

## Tissue Location


Detected in dendritic cells (at protein level). Highly expressed in the Reed-Sternberg cells of Hodgkin disease

## CLIP1 Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


## CLIP1 Antibody (N-term) - Images




All lanes: Anti-CLIP1 Antibody (N-term) at 1:1000 dilution + HL-60 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 162 KDa Blocking/Dilution buffer: 5% NFDm/TBST.




All lanes: Anti-CLIP1 Antibody (N-term) at 1:1000 dilution Lane 1: MCF-7 whole cell lysate Lane 2: HeLa whole cell lysate Lane 3: 293T whole cell lysate Lane 4: Rat brain lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 162 KDa Blocking/Dilution buffer: 5% NFDm/TBST.



Anti-CLIP1 Antibody (N-term) at 1:1000 dilution + A549 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 162 kDa Blocking/Dilution buffer: 5% NFDm/TBST.



CLIP1 Antibody (N-term) (Cat. #AP8950a) IHC analysis in formalin fixed and paraffin embedded mouse brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the CLIP1 Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.



CLIP1 Antibody (N-term) (Cat. #AP8950a) flow cytometric analysis of HeLa cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

#### **CLIP1 Antibody (N-term) - Background**

CLIP1 may be an intermediate filament associated protein that links endocytic vesicles to microtubules.

#### **CLIP1 Antibody (N-term) - References**

Yang, X., et al., J. Biol. Chem. 284 (42), 28775-28782 (2009)  
Meunier, B., et al., Eur. J. Cell Biol. 88 (2), 91-102 (2009)