

CLIP1 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP8950a

Specification

CLIP1 Antibody (N-term) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	P30622
Other Accession	O55156 , O9Z0H8 , O9UDT6 , O922J3 , O42184 , O9JK25
Reactivity	Human
Predicted	Chicken, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	162246
Antigen Region	228-254

CLIP1 Antibody (N-term) - Additional Information

Gene ID 6249

Other Names

CAP-Gly domain-containing linker protein 1, Cytoplasmic linker protein 1, Cytoplasmic linker protein 170 alpha-2, CLIP-170, Reed-Sternberg intermediate filament-associated protein, Restin, CLIP1, CYLN1, RSN

Target/Specificity

This CLIP1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 228-254 amino acids from the N-terminal region of human CLIP1.

Dilution

WB~~1:1000
IHC-P~~1:50~100
FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CLIP1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

CLIP1 Antibody (N-term) - Protein Information

Name CLIP1

Synonyms CYLN1, RSN

Function Binds to the plus end of microtubules and regulates the dynamics of the microtubule cytoskeleton. Promotes microtubule growth and microtubule bundling. Links cytoplasmic vesicles to microtubules and thereby plays an important role in intracellular vesicle trafficking. Plays a role in macropinocytosis and endosome trafficking.

Cellular Location

Cytoplasm. Cytoplasm, cytoskeleton. Cytoplasmic vesicle membrane; Peripheral membrane protein; Cytoplasmic side. Cell projection, ruffle. Note=Localizes to microtubule plus ends (PubMed:17889670, PubMed:21646404). Localizes preferentially to the ends of tyrosinated microtubules (By similarity). Accumulates in plasma membrane regions with ruffling and protrusions. Associates with the membranes of intermediate macropinocytic vesicles (PubMed:12433698) {ECO:0000250|UniProtKB:Q922J3, ECO:0000269|PubMed:12433698, ECO:0000269|PubMed:17889670, ECO:0000269|PubMed:21646404}

Tissue Location


Detected in dendritic cells (at protein level). Highly expressed in the Reed-Sternberg cells of Hodgkin disease

CLIP1 Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


CLIP1 Antibody (N-term) - Images




All lanes: Anti-CLIP1 Antibody (N-term) at 1:1000 dilution Lane 1: MCF-7 whole cell lysate Lane 2: Hela whole cell lysate Lane 3: 293T whole cell lysate Lane 4: Rat brain lysate Lysates/proteins at 20 µg per lane. Secondary: Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated (ASP1615) at 1/15000 dilution. Observed band size: 162 KDa Blocking/Dilution buffer: 5% NFDm/TBST.



Anti-CLIP1 Antibody (N-term) at 1:1000 dilution + A549 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 162 kDa Blocking/Dilution buffer: 5% NFDm/TBST.



CLIP1 Antibody (N-term) (Cat. #AP8950a) IHC analysis in formalin fixed and paraffin embedded mouse brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the CLIP1 Antibody (N-term) for immunohistochemistry. Clinical relevance has not been evaluated.



CLIP1 Antibody (N-term) - Background

CLIP1 may be an intermediate filament associated protein that links endocytic vesicles to microtubules.

CLIP1 Antibody (N-term) - References

Yang, X., et al., J. Biol. Chem. 284 (42), 28775-28782 (2009)
Meunier, B., et al., Eur. J. Cell Biol. 88 (2), 91-102 (2009)