

CCR2/CKR2 Antibody
Rabbit mAb
Catalog # AP90119**Specification****CCR2/CKR2 Antibody - Product Information**

| | |
|---|------------------------|
| Application | IHC, FC, IP |
| Primary Accession | P41597 |
| Clonality | Monoclonal |
| Other Names | |
| C-C chemokine receptor type 2; C-C CKR-2; CC chemokine receptor type 2; CC CKR 2; CC-CKR-2; CCCR2; CCR 2; CCR-2;; CCR1L;; CCR2; | |
| Isotype | Rabbit IgG |
| Host | Rabbit |
| Calculated MW | 41915 Da |

CCR2/CKR2 Antibody - Additional Information

| | |
|------------------------------|--|
| Purification | Affinity-chromatography |
| Immunogen | A synthesized peptide derived from human CCR2/CKR2 |
| Description | Receptor for the MCP-1, MCP-3 and MCP-4 chemokines. Transduces a signal by increasing the intracellular calcium ions level. Alternative coreceptor with CD4 for HIV-1 infection. |
| Storage Condition and Buffer | Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle. |

CCR2/CKR2 Antibody - Protein Information**Name** CCR2**Synonyms** CMKBR2**Function**

Key functional receptor for CCL2 but can also bind CCL7, and CCL12 (PubMed: [23408426](http://www.uniprot.org/citations/23408426), PubMed: [38157855](http://www.uniprot.org/citations/38157855), PubMed: [8048929](http://www.uniprot.org/citations/8048929), PubMed: [8146186](http://www.uniprot.org/citations/8146186)). Also transduces signaling mediated by CCL13 (PubMed: [38157855](http://www.uniprot.org/citations/38157855)). Its binding with CCL2 on monocytes and macrophages mediates chemotaxis and migration induction through the activation of the PI3K cascade, the small G protein Rac and lamellipodium protrusion (PubMed: [38157855](http://www.uniprot.org/citations/38157855)).

href="http://www.uniprot.org/citations/38157855" target="_blank">38157855). Also acts as a receptor for the beta-defensin DEFB106A/DEFB106B (PubMed:23938203). Regulates the expression of T-cell inflammatory cytokines and T-cell differentiation, promoting the differentiation of T-cells into T-helper 17 cells (Th17) during inflammation (By similarity). Facilitates the export of mature thymocytes by enhancing directional movement of thymocytes to sphingosine-1-phosphate stimulation and up-regulation of S1P1R expression; signals through the JAK-STAT pathway to regulate FOXO1 activity leading to an increased expression of S1P1R (By similarity). Plays an important role in mediating peripheral nerve injury-induced neuropathic pain (By similarity). Increases NMDA-mediated synaptic transmission in both dopamine D1 and D2 receptor-containing neurons, which may be caused by MAPK/ERK-dependent phosphorylation of GRIN2B/NMDAR2B (By similarity). Mediates the recruitment of macrophages and monocytes to the injury site following brain injury (By similarity).

Cellular Location

Cell membrane; Multi-pass membrane protein. Note=The chemoattractant receptors are distributed throughout the cell surface; after stimulation with a ligand, such as CCL2, they are rapidly recruited into microdomain clusters at the cell membrane.

Tissue Location

Expressed by monocytes and IL2-activated NK cells (PubMed:9058802). Abundantly expressed on CD14+/CD16- monocytes and weakly on CD14+/CD16+ monocytes, type 2 dendritic cells (DCs) and plasmacytoid DCs (at protein level) (PubMed:38157855)

CCR2/CKR2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CCR2/CKR2 Antibody - Images