

Histone H3 (mono methyl R17) Antibody
Rabbit mAb
Catalog # AP90268**Specification**

Histone H3 (mono methyl R17) Antibody - Product Information

Application	WB, IHC, ICC
Primary Accession	P68431
Reactivity	Rat
Clonality	Monoclonal
Other Names	
H3 histone; HIST1H3A; Histone cluster 1, H3a; H3R17me1	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	15404 Da

Histone H3 (mono methyl R17) Antibody - Additional Information

Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Histone H3 (mono methyl R17)
Description	H3 Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling. The nucleosome is a histone octamer containing two molecules each of H2A, H2B, H3 and H4 assembled in one H3-H4 heterotetramer and two H2A-H2B heterodimers.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Histone H3 (mono methyl R17) Antibody - Protein Information**Name** H3C1 ([HGNC:4766](#))**Synonyms** H3FA, HIST1H3A

Function

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.


Cellular Location

Nucleus. Chromosome.

Histone H3 (mono methyl R17) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Histone H3 (mono methyl R17) Antibody - Images

Western blot analysis of Histone H3 (mono methyl R17) expression in HeLa cell lysate.