

**Calreticulin Antibody**  
Rabbit mAb  
Catalog # AP90392

## Specification

---

### Calreticulin Antibody - Product Information

Application	WB, IHC, FC, ICC, IP
Primary Accession	<a href="#">P27797</a>
Reactivity	Rat
Clonality	Monoclonal
<b>Other Names</b>	
CALR; CRT; FLJ26680; RO; SSA; cC1Qr; CRP55; CRTC; HACBP;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	48142 Da

### Calreticulin Antibody - Additional Information

Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Calreticulin - ER Marker
Description	Calcium is a universal signaling molecule involved in many cellular functions such as cell motility, metabolism, protein modification, protein folding and apoptosis. Calcium is stored in the endoplasmic reticulum (ER), where it is buffered by calcium binding chaperones such as calnexin and calreticulin, and is released via the IP3 Receptor.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

### Calreticulin Antibody - Protein Information

Name CALR ([HGNC:1455](#))

Synonyms CRTC

#### Function

Calcium-binding chaperone that promotes folding, oligomeric assembly and quality control in the endoplasmic reticulum (ER) via the calreticulin/calnexin cycle. This lectin interacts transiently with almost all of the monoglucosylated glycoproteins that are synthesized in the ER (PubMed:<a href="http://www.uniprot.org/citations/7876246" target="\_blank">7876246</a>). Interacts with the DNA-binding domain of NR3C1 and mediates its nuclear export (PubMed:<a

href="http://www.uniprot.org/citations/11149926" target="\_blank">11149926</a>). Involved in maternal gene expression regulation. May participate in oocyte maturation via the regulation of calcium homeostasis (By similarity). Present in the cortical granules of non-activated oocytes, is exocytosed during the cortical reaction in response to oocyte activation and might participate in the block to polyspermy (By similarity).

#### Cellular Location


Endoplasmic reticulum lumen. Cytoplasm, cytosol. Secreted, extracellular space, extracellular matrix. Cell surface. Sarcoplasmic reticulum lumen {ECO:0000250|UniProtKB:P28491}. Cytoplasmic vesicle, secretory vesicle, Cortical granule {ECO:0000250|UniProtKB:Q8K3H7}. Cytolytic granule. Note=Also found in cell surface (T cells), cytosol and extracellular matrix (PubMed:10358038). During oocyte maturation and after parthenogenetic activation accumulates in cortical granules. In pronuclear and early cleaved embryos localizes weakly to cytoplasm around nucleus and more strongly in the region near the cortex (By similarity). In cortical granules of non-activated oocytes, is exocytosed during the cortical reaction in response to oocyte activation (By similarity). {ECO:0000250|UniProtKB:P28491, ECO:0000250|UniProtKB:Q8K3H7, ECO:0000269|PubMed:8418194}

#### Calreticulin Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Calreticulin Antibody - Images



Western blot analysis of Calreticulin expression in (1)HeLa cell lysate; (2)HepG2 cell lysate.