

**Cyclin D1 Antibody**  
Rabbit mAb  
Catalog # AP90436

## Specification

---

### Cyclin D1 Antibody - Product Information

Application	WB, IHC, ICC, IP
Primary Accession	<a href="#">P24385</a>
Reactivity	Rat
Clonality	Monoclonal
<b>Other Names</b>	
CCND1;BCL1; D11S287E; PRAD1; U21B31; Cyclin D1;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	33729 Da

### Cyclin D1 Antibody - Additional Information

Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Cyclin D1
Description	Phosphorylation of RB1 allows dissociation of the transcription factor E2F from the RB/E2F complex and the subsequent transcription of E2F target genes which are responsible for the progression through the G1 phase. Hypophosphorylates RB1 in early G1 phase. Cyclin D-CDK4 complexes are major integrators of various mitogenic and antimitogenic signals.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

### Cyclin D1 Antibody - Protein Information

**Name** CCND1 {ECO:0000303|PubMed:8204893, ECO:0000312|HGNC:HGNC:1582}

#### Function

Regulatory component of the cyclin D1-CDK4 (DC) complex that phosphorylates and inhibits members of the retinoblastoma (RB) protein family including RB1 and regulates the cell-cycle during G(1)/S transition (PubMed: [1827756](http://www.uniprot.org/citations/1827756), PubMed: [1833066](http://www.uniprot.org/citations/1833066), PubMed: [19412162](http://www.uniprot.org/citations/19412162), PubMed: [33854235](http://www.uniprot.org/citations/33854235), PubMed: [8114739](http://www.uniprot.org/citations/8114739))

target="\_blank">8114739</a>, PubMed:<a href="http://www.uniprot.org/citations/8302605" target="\_blank">8302605</a>). Phosphorylation of RB1 allows dissociation of the transcription factor E2F from the RB/E2F complex and the subsequent transcription of E2F target genes which are responsible for the progression through the G(1) phase (PubMed:<a href="http://www.uniprot.org/citations/1827756" target="\_blank">1827756</a>, PubMed:<a href="http://www.uniprot.org/citations/1833066" target="\_blank">1833066</a>, PubMed:<a href="http://www.uniprot.org/citations/19412162" target="\_blank">19412162</a>, PubMed:<a href="http://www.uniprot.org/citations/8114739" target="\_blank">8114739</a>, PubMed:<a href="http://www.uniprot.org/citations/8302605" target="\_blank">8302605</a>). Hypophosphorylates RB1 in early G(1) phase (PubMed:<a href="http://www.uniprot.org/citations/1827756" target="\_blank">1827756</a>, PubMed:<a href="http://www.uniprot.org/citations/1833066" target="\_blank">1833066</a>, PubMed:<a href="http://www.uniprot.org/citations/19412162" target="\_blank">19412162</a>, PubMed:<a href="http://www.uniprot.org/citations/8114739" target="\_blank">8114739</a>, PubMed:<a href="http://www.uniprot.org/citations/8302605" target="\_blank">8302605</a>). Cyclin D-CDK4 complexes are major integrators of various mitogenic and antimitogenic signals (PubMed:<a href="http://www.uniprot.org/citations/1827756" target="\_blank">1827756</a>, PubMed:<a href="http://www.uniprot.org/citations/1833066" target="\_blank">1833066</a>, PubMed:<a href="http://www.uniprot.org/citations/19412162" target="\_blank">19412162</a>, PubMed:<a href="http://www.uniprot.org/citations/8302605" target="\_blank">8302605</a>). Also a substrate for SMAD3, phosphorylating SMAD3 in a cell-cycle-dependent manner and repressing its transcriptional activity (PubMed:<a href="http://www.uniprot.org/citations/15241418" target="\_blank">15241418</a>). Component of the ternary complex, cyclin D1/CDK4/CDKN1B, required for nuclear translocation and activity of the cyclin D-CDK4 complex (PubMed:<a href="http://www.uniprot.org/citations/9106657" target="\_blank">9106657</a>). Exhibits transcriptional corepressor activity with INSM1 on the NEUROD1 and INS promoters in a cell cycle-independent manner (PubMed:<a href="http://www.uniprot.org/citations/16569215" target="\_blank">16569215</a>, PubMed:<a href="http://www.uniprot.org/citations/18417529" target="\_blank">18417529</a>).

#### Cellular Location


Nucleus. Cytoplasm Nucleus membrane. Note=Cyclin D-CDK4 complexes accumulate at the nuclear membrane and are then translocated to the nucleus through interaction with KIP/CIP family members


#### Cyclin D1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### Cyclin D1 Antibody - Images





Western blot analysis of Cyclin D1 expression in (1)MCF-7 cell lysates;(2) LnCaP cell lysates.