

**M6PR/IGF2R Antibody**  
Rabbit mAb  
Catalog # AP90519

**Specification**

---

**M6PR/IGF2R Antibody - Product Information**

Application	WB, IHC, FC, ICC, IP
Primary Accession	<a href="#">P11717</a>
Reactivity	Rat
Clonality	Monoclonal
<b>Other Names</b>	
IGF2R; CI Man-6-P receptor; CI-MPR; M6PR; MPR 300; Insulin-like growth factor 2 receptor; M6P/IGF2R; CD222;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	274375 Da

**M6PR/IGF2R Antibody - Additional Information**

Purification	<b>Affinity-chromatography</b>
Immunogen	<b>A synthesized peptide derived from human M6PR/IGF2R</b>
Description	<b>Transport of phosphorylated lysosomal enzymes from the Golgi complex and the cell surface to lysosomes. Lysosomal enzymes bearing phosphomannosyl residues bind specifically to mannose-6-phosphate receptors in the Golgi apparatus and the resulting receptor-ligand complex is transported to an acidic prelysosomal compartment where the low pH mediates the dissociation of the complex. This receptor also binds IGF2. Acts as a positive regulator of T-cell coactivation, by binding DPP4.</b>
Storage Condition and Buffer	<b>Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.</b>

**M6PR/IGF2R Antibody - Protein Information**

**Name** IGF2R  
**Synonyms** MPRI  
**Function**

Mediates the transport of phosphorylated lysosomal enzymes from the Golgi complex and the cell surface to lysosomes (PubMed:<a href="http://www.uniprot.org/citations/18817523" target="\_blank">18817523</a>, PubMed:<a href="http://www.uniprot.org/citations/2963003" target="\_blank">2963003</a>). Lysosomal enzymes bearing phosphomannosyl residues bind specifically to mannose-6-phosphate receptors in the Golgi apparatus and the resulting receptor-ligand complex is transported to an acidic prelysosomal compartment where the low pH mediates the dissociation of the complex (PubMed:<a href="http://www.uniprot.org/citations/18817523" target="\_blank">18817523</a>, PubMed:<a href="http://www.uniprot.org/citations/2963003" target="\_blank">2963003</a>). The receptor is then recycled back to the Golgi for another round of trafficking through its binding to the retromer (PubMed:<a href="http://www.uniprot.org/citations/18817523" target="\_blank">18817523</a>). This receptor also binds IGF2 (PubMed:<a href="http://www.uniprot.org/citations/18046459" target="\_blank">18046459</a>). Acts as a positive regulator of T-cell coactivation by binding DPP4 (PubMed:<a href="http://www.uniprot.org/citations/10900005" target="\_blank">10900005</a>).

### Cellular Location


Golgi apparatus membrane; Single-pass type I membrane protein. Endosome membrane; Single-pass type I membrane protein. Note=Mainly localized in the Golgi at steady state and not detectable in lysosome (PubMed:18817523) Colocalized with DPP4 in internalized cytoplasmic vesicles adjacent to the cell surface (PubMed:10900005).

### M6PR/IGF2R Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### M6PR/IGF2R Antibody - Images



Western blot analysis of extracts of M6PR expression in Jurkat cell lysate.