

Aurora A Antibody
Rabbit mAb
Catalog # AP90525

Specification

Aurora A Antibody - Product Information

Application	WB, FC, ICC
Primary Accession	O14965
Clonality	Monoclonal
Other Names	
AIK, ARK1, AYK1, Aurora-A, Aurora-related kinase 1, BTAK, IAK1, Ipl1- and aurora-related kinase 1, STK15, STK6, Serine/threonine kinase 15	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	45809 Da

Aurora A Antibody - Additional Information

Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Aurora A
Description	Mitotic serine/threonine kinases that contributes to the regulation of cell cycle progression. Associates with the centrosome and the spindle microtubules during mitosis and plays a critical role in various mitotic events including the establishment of mitotic spindle, centrosome duplication, centrosome separation as well as maturation, chromosomal alignment, spindle assembly checkpoint, and cytokinesis.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Aurora A Antibody - Protein Information

Name AURKA ([HGNC:11393](#))

Function

Mitotic serine/threonine kinase that contributes to the regulation of cell cycle progression
(PubMed: [11039908](http://www.uniprot.org/citations/11039908)),
(PubMed: [12390251](http://www.uniprot.org/citations/12390251)),
(PubMed: [17125279](http://www.uniprot.org/citations/17125279)),
(PubMed: [17360485](http://www.uniprot.org/citations/17360485)),

PubMed: 18615013, PubMed: 26246606). Associates with the centrosome and the spindle microtubules during mitosis and plays a critical role in various mitotic events including the establishment of mitotic spindle, centrosome duplication, centrosome separation as well as maturation, chromosomal alignment, spindle assembly checkpoint, and cytokinesis (PubMed: 14523000, PubMed: 26246606). Required for normal spindle positioning during mitosis and for the localization of NUMA1 and DCTN1 to the cell cortex during metaphase (PubMed: 27335426). Required for initial activation of CDK1 at centrosomes (PubMed: 13678582, PubMed: 15128871). Phosphorylates numerous target proteins, including ARHGEF2, BORA, BRCA1, CDC25B, DLGP5, HDAC6, KIF2A, LATS2, NDEL1, PARD3, PPP1R2, PLK1, RASSF1, TACC3, p53/TP53 and TPX2 (PubMed: 11551964, PubMed: 14702041, PubMed: 15128871, PubMed: 15147269, PubMed: 15987997, PubMed: 17604723, PubMed: 18056443, PubMed: 18615013). Regulates KIF2A tubulin depolymerase activity (PubMed: 19351716). Important for microtubule formation and/or stabilization (PubMed: 18056443). Required for normal axon formation (PubMed: 19812038). Plays a role in microtubule remodeling during neurite extension (PubMed: 19668197). Also acts as a key regulatory component of the p53/TP53 pathway, and particularly the checkpoint- response pathways critical for oncogenic transformation of cells, by phosphorylating and destabilizing p53/TP53 (PubMed: 14702041). Phosphorylates its own inhibitors, the protein phosphatase type 1 (PP1) isoforms, to inhibit their activity (PubMed: 11551964). Inhibits cilia outgrowth (By similarity). Required for cilia disassembly via phosphorylation of HDAC6 and subsequent deacetylation of alpha-tubulin (PubMed: 17604723, PubMed: 20643351). Regulates protein levels of the anti-apoptosis protein BIRC5 by suppressing the expression of the SCF(FBXL7) E3 ubiquitin-protein ligase substrate adapter FBXL7 through the phosphorylation of the transcription factor FOXP1 (PubMed: 28218735).

Cellular Location

Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle pole. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriole {ECO:0000250|UniProtKB:P97477}. Cell projection, neuron projection {ECO:0000250|UniProtKB:P97477}. Cell projection, cilium. Cytoplasm, cytoskeleton, cilium basal body. Basolateral cell membrane {ECO:0000250|UniProtKB:F1PNY0}. Note=Detected at the neurite hillock in developing neurons (By similarity). Localizes at the centrosome in mitotic cells from early prophase until telophase, but also localizes to the spindle pole MTs from prophase to anaphase (PubMed:17229885, PubMed:21225229, PubMed:9606188). Colocalized with SIRT2 at centrosome (PubMed:22014574). Moves to the midbody during both telophase and cytokinesis (PubMed:17726514). Associates with both the pericentriolar material (PCM) and centrioles (PubMed:22014574). The localization to the spindle poles is regulated by AAAS

(PubMed:26246606) {ECO:0000250|UniProtKB:P97477, ECO:0000269|PubMed:17229885, ECO:0000269|PubMed:17726514, ECO:0000269|PubMed:21225229, ECO:0000269|PubMed:22014574, ECO:0000269|PubMed:26246606, ECO:0000269|PubMed:9606188}

Tissue Location


Highly expressed in testis and weakly in skeletal muscle, thymus and spleen. Also highly expressed in colon, ovarian, prostate, neuroblastoma, breast and cervical cancer cell lines

Aurora A Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Aurora A Antibody - Images



Western blot analysis of Aurora A expression in HepG2 cell lysate.