

**Prion Protein Antibody**  
**Rabbit mAb**  
**Catalog # AP90618****Specification**

---

**Prion Protein Antibody - Product Information**

Application	WB, IHC, FC, ICC
Primary Accession	<a href="#">P04156</a>
Reactivity	Rat
Clonality	Monoclonal
<b>Other Names</b>	
CJD; GSS; PrP; PRNP; ASCR; KURU; PRIP; PrPc; CD230; AltPrP; Prion protein; PrP27-30; PrP33-35C;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	27661 Da

**Prion Protein Antibody - Additional Information**

Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Prion Protein
Description	The PRNP gene encodes the major prion protein (PrP, CD230), a widely-expressed glycoprotein expressed at high levels in the central nervous system. While the typical cellular function of PrP is not well defined, it is a putative antioxidant and a metal-binding protein that may be involved in signal transduction.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

**Prion Protein Antibody - Protein Information****Name** PRNP**Synonyms** ALTPRP, PRIP, PRP**Function**

Its primary physiological function is unclear. May play a role in neuronal development and synaptic plasticity. May be required for neuronal myelin sheath maintenance. May promote myelin homeostasis through acting as an agonist for ADGRG6 receptor. May play a role in iron uptake and iron homeostasis. Soluble oligomers are toxic to cultured neuroblastoma cells and induce apoptosis (in vitro) (By similarity). Association with GPC1 (via its heparan sulfate chains) targets PRNP to lipid rafts. Also provides Cu(2+) or Zn(2+) for the ascorbate-mediated GPC1 deaminase

degradation of its heparan sulfate side chains (By similarity).

#### **Cellular Location**

Cell membrane; Lipid-anchor, GPI-anchor. Golgi apparatus {ECO:0000250|UniProtKB:P04925}.


Note=Targeted to lipid rafts via association with the heparan sulfate chains of GPC1. Colocalizes, in the presence of Cu(2+), to vesicles in para- and perinuclear regions, where both proteins undergo internalization. Heparin displaces PRNP from lipid rafts and promotes endocytosis.

#### **Prion Protein Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **Prion Protein Antibody - Images**



Western blot analysis of Prion Protein expression in Hunam fetal brain lysate.