

PRP19 Antibody
Rabbit mAb
Catalog # AP90683

Specification

PRP19 Antibody - Product Information

Application	WB, IHC, FC, ICC
Primary Accession	O9UMS4
Reactivity	Rat
Clonality	Monoclonal

Other Names

NMP200; nuclear matrix protein 200; pre-mRNA-processing factor 19; PRP19; PRP19/PSO4; PRPF19; PSO4; senescence evasion factor; SNEV; UBOX4;

Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	55181 Da

PRP19 Antibody - Additional Information

Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human PRP19

Description	Ubiquitin-protein ligase which is a core component of several complexes mainly involved pre-mRNA splicing and DNA repair. Core component of the PRP19C/Prp19 complex/NTC/Nineteen complex which is part of the spliceosome and participates in its assembly, its remodeling and is required for its activity. During assembly of the spliceosome, mediates 'Lys-63'-linked polyubiquitination of the U4 spliceosomal protein PRPF3. Ubiquitination of PRPF3 allows its recognition by the U5 component PRPF8 and stabilizes the U4/U5/U6 tri-snRNP spliceosomal complex (PubMed:20595234).
-------------	--

Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
------------------------------	---

PRP19 Antibody - Protein Information

Name PRPF19 ([HGNC:17896](#))

Function

Ubiquitin-protein ligase which is a core component of several complexes mainly involved pre-mRNA splicing and DNA repair. Required for pre-mRNA splicing as component of the spliceosome (PubMed: [28076346](http://www.uniprot.org/citations/28076346)), PubMed: [28502770](http://www.uniprot.org/citations/28502770), PubMed: [29301961](http://www.uniprot.org/citations/29301961), PubMed: [29360106](http://www.uniprot.org/citations/29360106), PubMed: [30705154](http://www.uniprot.org/citations/30705154)). Core component of the PRP19C/Prp19 complex/NTC/Nineteen complex which is part of the spliceosome and participates in its assembly, its remodeling and is required for its activity. During assembly of the spliceosome, mediates 'Lys-63'-linked polyubiquitination of the U4 spliceosomal protein PRPF3. Ubiquitination of PRPF3 allows its recognition by the U5 component PRPF8 and stabilizes the U4/U5/U6 tri- snRNP spliceosomal complex (PubMed: [20595234](http://www.uniprot.org/citations/20595234)). Recruited to RNA polymerase II C-terminal domain (CTD) and the pre-mRNA, it may also couple the transcriptional and spliceosomal machineries (PubMed: [21536736](http://www.uniprot.org/citations/21536736)). The XAB2 complex, which contains PRPF19, is also involved in pre-mRNA splicing, transcription and transcription-coupled repair (PubMed: [17981804](http://www.uniprot.org/citations/17981804)). Beside its role in pre-mRNA splicing PRPF19, as part of the PRP19-CDC5L complex, plays a role in the DNA damage response/DDR. It is recruited to the sites of DNA damage by the RPA complex where PRPF19 directly ubiquitinates RPA1 and RPA2. 'Lys-63'-linked polyubiquitination of the RPA complex allows the recruitment of the ATR-ATRIP complex and the activation of ATR, a master regulator of the DNA damage response (PubMed: [24332808](http://www.uniprot.org/citations/24332808)). May also play a role in DNA double-strand break (DSB) repair by recruiting the repair factor SETMAR to altered DNA (PubMed: [18263876](http://www.uniprot.org/citations/18263876)). As part of the PSO4 complex may also be involved in the DNA interstrand cross-links/ICLs repair process (PubMed: [16223718](http://www.uniprot.org/citations/16223718)). In addition, may also mediate 'Lys-48'-linked polyubiquitination of substrates and play a role in proteasomal degradation (PubMed: [11435423](http://www.uniprot.org/citations/11435423)). May play a role in the biogenesis of lipid droplets (By similarity). May play a role in neural differentiation possibly through its function as part of the spliceosome (By similarity).

Cellular Location

Nucleus. Nucleus, nucleoplasm. Cytoplasm, cytoskeleton, spindle. Cytoplasm. Lipid droplet {ECO:0000250|UniProtKB:Q99KP6}. Note=Nucleoplasmic in interphase cells Irregularly distributed in anaphase cells. In prophase cells, uniformly distributed, but not associated with condensing chromosomes. Found in extrachromosomal regions in metaphase cells. Mainly localized to the mitotic spindle apparatus when chromosomes segregate during anaphase When nuclei reform during late telophase, uniformly distributed in daughter cells and displays no preferred association with decondensing chromatin. Recruited on damaged DNA at sites of double-strand break

Tissue Location

Ubiquitous. Weakly expressed in senescent cells of different tissue origins. Highly expressed in tumor cell lines


PRP19 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PRP19 Antibody - Images



Western blot analysis of PRP19 expression in (1) HepG2 cell lysate; (2) NIH/3T3 cell lysate.