

**MAP1LC3A Antibody**  
Rabbit mAb  
Catalog # AP90771**Specification****MAP1LC3A Antibody - Product Information**

Application	WB, IHC, ICC, IP
Primary Accession	<a href="#">O9H492</a>
Reactivity	Rat
Clonality	Monoclonal

**Other Names**

ATG8E; Autophagy-related ubiquitin-like modifier LC3 A; LC3; LC3A; MAP1 light chain 3 like protein 1; MAP1A/1B light chain 3A;

Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	14272 Da

**MAP1LC3A Antibody - Additional Information**

Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human MAP1LC3A
Description	Autophagy is generally activated by conditions of nutrient deprivation, but it has also been associated with a number of physiological processes including development, differentiation, neurodegenerative diseases, infection, and cancer. The presence of LC3 in autophagosomes and the conversion of LC3 to the lower migrating form, LC3-II, have been used as indicators of autophagy.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

**MAP1LC3A Antibody - Protein Information**

**Name** MAP1LC3A

**Function**

Ubiquitin-like modifier involved in formation of autophagosomal vacuoles (autophagosomes) (PubMed:<a href="http://www.uniprot.org/citations/20713600" target="\_blank">20713600</a>, PubMed:<a href="http://www.uniprot.org/citations/24290141" target="\_blank">24290141</a>). While LC3s are involved in elongation of the phagophore membrane, the GABARAP/GATE-16

subfamily is essential for a later stage in autophagosome maturation (PubMed:<a href="http://www.uniprot.org/citations/20713600" target="\_blank">20713600</a>). Through its interaction with the reticulophagy receptor TEX264, participates in the remodeling of subdomains of the endoplasmic reticulum into autophagosomes upon nutrient stress, which then fuse with lysosomes for endoplasmic reticulum turnover (PubMed:<a href="http://www.uniprot.org/citations/31006537" target="\_blank">31006537</a>, PubMed:<a href="http://www.uniprot.org/citations/31006538" target="\_blank">31006538</a>).

#### Cellular Location

Cytoplasmic vesicle, autophagosome membrane; Lipid-anchor. Endomembrane system; Lipid-anchor. Cytoplasm, cytoskeleton {ECO:0000250|UniProtKB:Q91VR7}. Note=LC3-II binds to the autophagic membranes.

#### Tissue Location


Most abundant in heart, brain, liver, skeletal muscle and testis but absent in thymus and peripheral blood leukocytes

### MAP1LC3A Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### MAP1LC3A Antibody - Images



Western blot analysis of MAP1LC3A expression in Human brain lysate.