

RAB7 Antibody
Rabbit mAb
Catalog # AP90815

Specification

RAB7 Antibody - Product Information

Application	WB, IHC, FC, ICC
Primary Accession	P51149
Reactivity	Rat
Clonality	Monoclonal

Other Names

CMT2B; PRO2706; PSN; RAB7, member RAS oncogene family; RAB7A; RAB7A, member RAS oncogene family; Ras associated protein RAB7; Ras related protein Rab7a;

Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	23490 Da

RAB7 Antibody - Additional Information

Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human RAB7
Description	Plays a role in the fusion of phagosomes with lysosomes. Plays important roles in microbial pathogen infection and survival, as well as in participating in the life cycle of viruses. Microbial pathogens possess survival strategies governed by RAB7A, sometimes by employing RAB7A function (e.g. Salmonella) and sometimes by excluding RAB7A function.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

RAB7 Antibody - Protein Information

Name RAB7A ([HGNC:9788](#))

Synonyms RAB7

Function

Small GTPase which cycles between active GTP-bound and inactive GDP-bound states. In its active state, binds to a variety of effector proteins playing a key role in the regulation of endo- lysosomal trafficking. Governs early-to-late endosomal maturation, microtubule minus-end as well as plus-end directed endosomal migration and positioning, and endosome-lysosome transport

through different protein-protein interaction cascades. Plays a central role, not only in endosomal traffic, but also in many other cellular and physiological events, such as growth-factor-mediated cell signaling, nutrient-transporter mediated nutrient uptake, neurotrophin transport in the axons of neurons and lipid metabolism. Also involved in regulation of some specialized endosomal membrane trafficking, such as maturation of melanosomes, pathogen-induced phagosomes (or vacuoles) and autophagosomes. Plays a role in the maturation and acidification of phagosomes that engulf pathogens, such as *S.aureus* and *M.tuberculosis*. Plays a role in the fusion of phagosomes with lysosomes. Plays important roles in microbial pathogen infection and survival, as well as in participating in the life cycle of viruses. Microbial pathogens possess survival strategies governed by RAB7A, sometimes by employing RAB7A function (e.g. *Salmonella*) and sometimes by excluding RAB7A function (e.g. *Mycobacterium*). In concert with RAC1, plays a role in regulating the formation of RBs (ruffled borders) in osteoclasts. Controls the endosomal trafficking and neurite outgrowth signaling of NTRK1/TRKA (PubMed:11179213, PubMed:12944476, PubMed:14617358, PubMed:20028791, PubMed:21255211). Regulates the endocytic trafficking of the EGF-EGFR complex by regulating its lysosomal degradation. Involved in the ADRB2-stimulated lipolysis through lipophagy, a cytosolic lipase-independent autophagic pathway (By similarity). Required for the exosomal release of SDCBP, CD63 and syndecan (PubMed:22660413). Required for vesicular trafficking and cell surface expression of ACE2 (PubMed:33147445). May play a role in PRPH neuronal intermediate filament assembly (By similarity).

Cellular Location

Cytoplasmic vesicle, phagosome membrane; Peripheral membrane protein; Cytoplasmic side. Late endosome membrane; Peripheral membrane protein; Cytoplasmic side. Lysosome membrane; Peripheral membrane protein; Cytoplasmic side. Melanosome membrane; Peripheral membrane protein; Cytoplasmic side. Cytoplasmic vesicle, autophagosome membrane; Peripheral membrane protein; Cytoplasmic side. Lipid droplet {ECO:0000250|UniProtKB:P51150}. Endosome membrane; Peripheral membrane protein. Cytoplasmic vesicle {ECO:0000250|UniProtKB:P51150} Mitochondrion membrane; Peripheral membrane protein. Note=Colocalizes with OSBPL1A at the late endosome (PubMed:16176980). Found in the ruffled border (a late endosomal-like compartment in the plasma membrane) of bone-resorbing osteoclasts. Recruited to phagosomes containing *S.aureus* or *Mycobacterium* (PubMed:21255211). Lipid droplet localization is increased upon ADRB2 stimulation (By similarity). Recruited to damaged mitochondria during mitophagy in a RIMOC1-dependent manner (PubMed:34432599). {ECO:0000250|UniProtKB:P51150, ECO:0000269|PubMed:16176980, ECO:0000269|PubMed:21255211, ECO:0000269|PubMed:34432599}

Tissue Location

Widely expressed; high expression found in skeletal muscle.


RAB7 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)

- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RAB7 Antibody - Images



Western blot analysis of RAB7 expression in (1) A375 cell lysate; (2) 3T3 cell lysate.