

MEK2 Antibody

Rabbit mAb Catalog # AP90863

Specification

MEK2 Antibody - Product Information

Application WB, IHC, FC, ICC, IP

Primary Accession P36507
Clonality Monoclonal

Other Names

Dual specificity mitogen-activated protein kinase kinase 2; MAP kinase kinase 2; MAPKK 2; ERK activator kinase 2; MAPK/ERK kinase 2; MEK 2; MEK-2; MEK-2; MEK-2; PRKMK2;

Isotype Rabbit IgG
Host Rabbit
Calculated MW 44424 Da

MEK2 Antibody - Additional Information

Purification Affinity-chromatography

Immunogen A synthesized peptide derived from human

MEK2

Description Catalyzes the concomitant phosphorylation

of a threonine and a tyrosine residue in a Thr-Glu-Tyr sequence located in MAP kinases. Activates the ERK1 and ERK2 MAP

kinases.

Storage Condition and Buffer Rabbit IgG in phosphate buffered saline,

pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid

freeze / thaw cycle.

MEK2 Antibody - Protein Information

Name MAP2K2

Synonyms MEK2, MKK2, PRKMK2

Function

Catalyzes the concomitant phosphorylation of a threonine and a tyrosine residue in a Thr-Glu-Tyr sequence located in MAP kinases. Activates the ERK1 and ERK2 MAP kinases (By similarity). Activates BRAF in a KSR1 or KSR2-dependent manner; by binding to KSR1 or KSR2 releases the inhibitory intramolecular interaction between KSR1 or KSR2 protein kinase and N-terminal domains which promotes KSR1 or KSR2-BRAF dimerization and BRAF activation (PubMed:29433126).

Cellular Location

Cytoplasm. Membrane; Peripheral membrane protein. Note=Membrane localization is probably




regulated by its interaction with KSR1.

MEK2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

MEK2 Antibody - Images



Western blot analysis of MEK2 expression in Jurkat cell lysate.