

HDAC9 Antibody
Rabbit mAb
Catalog # AP90890

Specification

HDAC9 Antibody - Product Information

Application	WB, IHC, ICC, IP
Primary Accession	O9UKV0
Reactivity	Rat
Clonality	Monoclonal

Other Names

HD9; Histone deacetylase 7B; HD7; HD7b; Histone deacetylase-related protein; HDAC9; HDAC7; HDAC7B; HDRP; MITR;

Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	111297 Da

HDAC9 Antibody - Additional Information

Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human HDAC9
Description	Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Represses MEF2-dependent transcription.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

HDAC9 Antibody - Protein Information

Name HDAC9

Synonyms HDAC7, HDAC7B, HDRP, KIAA0744, MITR

Function

Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Represses MEF2-dependent transcription.

Cellular Location

Nucleus.

Tissue Location


Broadly expressed, with highest levels in brain, heart, muscle and testis. Isoform 3 is present in human bladder carcinoma cells (at protein level).

HDAC9 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HDAC9 Antibody - Images



Western blot analysis of HDAC9 expression in K562 cell lysate.