

Phospho-Histone H2A.X (S139) Antibody

Rabbit mAb Catalog # AP91004

Specification

Phospho-Histone H2A.X (\$139) Antibody - Product Information

Application WB, IHC, ICC, IP

Primary Accession P16104
Reactivity Rat

Clonality Monoclonal

Other Names

Description

H2A.X; H2AFX; H2a/x; HIST5-2AX; Histone H2A.X;

Isotype Rabbit IgG
Host Rabbit
Calculated MW 15145 Da

Phospho-Histone H2A.X (S139) Antibody - Additional Information

Purification Affinity-chromatography

Immunogen A synthesized peptide derived from human

Phospho-Histone H2A.X (S139)
Variant histone H2A which replaces
conventional H2A in a subset of
nucleosomes. Nucleosomes wrap and
compact DNA into chromatin, limiting DNA
accessibility to the cellular machineries
which require DNA as a template. Histones
thereby play a central role in transcription

regulation, DNA repair, DNA replication and chromosomal stability.

Storage Condition and Buffer Rabbit IgG in phosphate buffered saline ,

pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid

freeze / thaw cycle.

Phospho-Histone H2A.X (S139) Antibody - Protein Information

Name H2AX (HGNC:4739)

Function

Variant histone H2A which replaces conventional H2A in a subset of nucleosomes. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post- translational modifications of histones, also called histone code, and nucleosome remodeling. Required for checkpoint-mediated arrest of cell cycle progression in response to low doses of ionizing radiation and for efficient repair of DNA double strand breaks (DSBs) specifically



when modified by C-terminal phosphorylation.


Cellular LocationNucleus. Chromosome

Phospho-Histone H2A.X (S139) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Phospho-Histone H2A.X (S139) Antibody - Images



Western blot analysis of HistoneH2A.X phosphorylation expression in Jurkat cell lysate treated with etoposide.