

HMGCR Antibody
Rabbit mAb
Catalog # AP91071

Specification

HMGCR Antibody - Product Information

Application	WB, IP
Primary Accession	P04035
Reactivity	Rat
Clonality	Monoclonal
Other Names	
3H3M; HMDH; HMG CoA reductase; HMG CoAR;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	97476 Da

HMGCR Antibody - Additional Information

Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human HMGCR
Description	Transmembrane glycoprotein that is the rate-limiting enzyme in cholesterol biosynthesis as well as in the biosynthesis of nonsterol isoprenoids that are essential for normal cell function including ubiquinone and geranylgeranyl proteins.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

HMGCR Antibody - Protein Information

Name HMGCR ([HGNC:5006](#))

Function

Catalyzes the conversion of (3S)-hydroxy-3-methylglutaryl-CoA (HMG-CoA) to mevalonic acid, the rate-limiting step in the synthesis of cholesterol and other isoprenoids, thus plays a critical role in cellular cholesterol homeostasis (PubMed: [21357570](http://www.uniprot.org/citations/21357570), PubMed: [2991281](http://www.uniprot.org/citations/2991281), PubMed: [36745799](http://www.uniprot.org/citations/36745799), PubMed: [6995544](http://www.uniprot.org/citations/6995544)). HMGCR is the main target of statins, a class of cholesterol-lowering drugs (PubMed: [11349148](http://www.uniprot.org/citations/11349148), PubMed: [18540668](http://www.uniprot.org/citations/18540668), PubMed: [36745799](http://www.uniprot.org/citations/36745799)).

target="_blank">36745799).

Cellular Location

Endoplasmic reticulum membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:P00347}. Peroxisome membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:P00347}

Tissue Location


[Isoform 1]: Ubiquitously expressed with the highest levels in the cerebellum, fetal brain, testis, skin and adrenal gland. [Isoform 3]: Low abundance except in skin, esophagus, and uterine cervix.

HMGR Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HMGR Antibody - Images



Western blot analysis of HMGR expression in Jurkat cell lysate.