

FGF2 Antibody
Rabbit mAb
Catalog # AP91122

Specification

FGF2 Antibody - Product Information

Application	WB, FC, IP
Primary Accession	P09038
Clonality	Monoclonal
Other Names	
Basic fibroblast growth factor; BFGF; FGF2; FGF2 basic; FGF2; FGFB; Fibroblast growth factor 2 (basic); HBGF2;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	30770 Da

FGF2 Antibody - Additional Information

Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human FGF2
Description	Plays an important role in the regulation of cell survival, cell division, angiogenesis, cell differentiation and cell migration. Functions as potent mitogen in vitro. Can induce angiogenesis.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

FGF2 Antibody - Protein Information

Name FGF2

Synonyms FGFB

Function

Acts as a ligand for FGFR1, FGFR2, FGFR3 and FGFR4 (PubMed:[8663044](http://www.uniprot.org/citations/8663044)). Also acts as an integrin ligand which is required for FGF2 signaling (PubMed:[28302677](http://www.uniprot.org/citations/28302677)). Binds to integrin ITGAV:ITGB3 (PubMed:[28302677](http://www.uniprot.org/citations/28302677)). Plays an important role in the regulation of cell survival, cell division, cell differentiation and cell migration (PubMed:[28302677](http://www.uniprot.org/citations/28302677), PubMed:[8663044](http://www.uniprot.org/citations/8663044)). Functions as a

potent mitogen in vitro (PubMed:1721615, PubMed:3732516, PubMed:3964259). Can induce angiogenesis (PubMed:23469107, PubMed:28302677). Mediates phosphorylation of ERK1/2 and thereby promotes retinal lens fiber differentiation (PubMed:29501879).

Cellular Location

Secreted. Nucleus. Note=Exported from cells by an endoplasmic reticulum (ER)/Golgi-independent mechanism. Unconventional secretion of FGF2 occurs by direct translocation across the plasma membrane (PubMed:20230531). Binding of exogenous FGF2 to FGFR facilitates endocytosis followed by translocation of FGF2 across endosomal membrane into the cytosol (PubMed:22321063). Nuclear import from the cytosol requires the classical nuclear import machinery, involving proteins KPNA1 and KPNB1, as well as CEP57 (PubMed:22321063)

Tissue Location


Expressed in granulosa and cumulus cells. Expressed in hepatocellular carcinoma cells, but not in non-cancerous liver tissue.

FGF2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FGF2 Antibody - Images



Western blot analysis of FGF2 expression in K562 cell lysate.