

**XPB1 Antibody**  
Rabbit mAb  
Catalog # AP91156

## Specification

---

### XPB1 Antibody - Product Information

|                    |                        |
|--------------------|------------------------|
| Application        | WB, IHC, FC, ICC       |
| Primary Accession  | <a href="#">P17861</a> |
| Clonality          | Monoclonal             |
| <b>Other Names</b> |                        |
| TREB5; XPB1; XBP2; |                        |
| Isotype            | Rabbit IgG             |
| Host               | Rabbit                 |
| Calculated MW      | 28695 Da               |

### XPB1 Antibody - Additional Information

|                              |   |
|------------------------------|---|
| Purification                 | <b>Affinity-chromatography</b>  |
| Immunogen                    | <b>A synthesized peptide derived from human XPB1</b>  |
| Description                  | <b>Acts during endoplasmic reticulum stress (ER) by activating unfolded protein response (UPR) target genes via direct binding to the UPR element (UPRE). Binds DNA preferably to the CRE-like element 5'-GATGACGTG[TG]N(3)[AT]T-3', and also to some TPA response elements (TRE). Binds to the HLA DR-alpha promoter. Binds to the Tax-responsive element (TRE) of HTLV-I.</b> |
| Storage Condition and Buffer | <b>Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.</b>  |

### XPB1 Antibody - Protein Information

**Name** XPB1 ([HGNC:12801](#))

#### Function

Functions as a transcription factor during endoplasmic reticulum (ER) stress by regulating the unfolded protein response (UPR). Required for cardiac myogenesis and hepatogenesis during embryonic development, and the development of secretory tissues such as exocrine pancreas and salivary gland (By similarity). Involved in terminal differentiation of B lymphocytes to plasma cells and production of immunoglobulins (PubMed:<a href="http://www.uniprot.org/citations/11460154" target="\_blank">11460154</a>). Modulates the cellular response to ER stress in a PIK3R-dependent manner (PubMed:<a href="http://www.uniprot.org/citations/20348923" target="\_blank">20348923</a>).

target="\_blank">20348923</a>). Binds to the cis-acting X box present in the promoter regions of major histocompatibility complex class II genes (PubMed:<a href="http://www.uniprot.org/citations/8349596" target="\_blank">8349596</a>). Involved in VEGF-induced endothelial cell (EC) proliferation and retinal blood vessel formation during embryonic development but also for angiogenesis in adult tissues under ischemic conditions. Functions also as a major regulator of the UPR in obesity-induced insulin resistance and type 2 diabetes for the management of obesity and diabetes prevention (By similarity).

#### Cellular Location

Endoplasmic reticulum. Note=Colocalizes with ERN1 and KDR in the endoplasmic reticulum in endothelial cells in a vascular endothelial growth factor (VEGF)-dependent manner (PubMed:23529610) [Isoform 2]: Nucleus. Cytoplasm {ECO:0000250|UniProtKB:O35426}. Note=Localizes predominantly in the nucleus. Colocalizes in the nucleus with SIRT1. Translocates into the nucleus in a PIK3R-, ER stress-induced- and/or insulin-dependent manner (By similarity). {ECO:0000250|UniProtKB:O35426}

#### Tissue Location


Expressed in plasma cells in rheumatoid synovium (PubMed:11460154). Over-expressed in primary breast cancer and metastatic breast cancer cells (PubMed:25280941). Isoform 1 and isoform 2 are expressed at higher level in proliferating as compared to confluent quiescent endothelial cells (PubMed:19416856)

### XBP1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### XBP1 Antibody - Images



Western blot analysis of XBP1 expression in Jurkat cell lysate.