

Lin28 Antibody
Rabbit mAb
Catalog # AP91157

Specification

Lin28 Antibody - Product Information

Application	WB, FC, IP
Primary Accession	O9H9Z2
Clonality	Monoclonal
Other Names	
Protein lin-28 homolog A; LIN28A; CSDD1; LIN28; ZCCHC1;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	22743 Da

Lin28 Antibody - Additional Information

Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Lin28
Description	Involved in the timing of developmental events and choice of stage specific cell fates. Acts as a suppressor of microRNA (miRNA) biogenesis by specifically binding the precursor let-7 (pre-let-7), a miRNA precursor. Acts by binding pre-let-7 and recruiting ZCCHC11/TUT4 uridylyltransferase, leading to the terminal uridylation of pre-let-7.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Lin28 Antibody - Protein Information

Name LIN28A

Synonyms CSDD1, LIN28, ZCCHC1

Function

RNA-binding protein that inhibits processing of pre-let-7 miRNAs and regulates translation of mRNAs that control developmental timing, pluripotency and metabolism (PubMed:21247876). Seems to recognize a common structural G-quartet (G4) feature in its miRNA and mRNA targets (Probable). 'Translational enhancer' that drives specific mRNAs to polysomes and increases the efficiency of protein synthesis. Its association with the translational machinery and target mRNAs results in an

increased number of initiation events per molecule of mRNA and, indirectly, in mRNA stabilization. Binds IGF2 mRNA, MYOD1 mRNA, ARBP/36B4 ribosomal protein mRNA and its own mRNA. Essential for skeletal muscle differentiation program through the translational up-regulation of IGF2 expression. Suppressor of microRNA (miRNA) biogenesis, including that of let-7, miR107, miR-143 and miR-200c. Specifically binds the miRNA precursors (pre-miRNAs), recognizing an 5'-GGAG-3' motif found in pre-miRNA terminal loop, and recruits TUT4 and TUT7 uridylyltransferases (PubMed: [18951094](http://www.uniprot.org/citations/18951094) target=" _blank">18951094, PubMed: [19703396](http://www.uniprot.org/citations/19703396) target=" _blank">19703396, PubMed: [22118463](http://www.uniprot.org/citations/22118463) target=" _blank">22118463, PubMed: [22898984](http://www.uniprot.org/citations/22898984) target=" _blank">22898984). This results in the terminal uridylation of target pre-miRNAs (PubMed: [18951094](http://www.uniprot.org/citations/18951094) target=" _blank">18951094, PubMed: [19703396](http://www.uniprot.org/citations/19703396) target=" _blank">19703396, PubMed: [22118463](http://www.uniprot.org/citations/22118463) target=" _blank">22118463, PubMed: [22898984](http://www.uniprot.org/citations/22898984) target=" _blank">22898984). Uridylated pre-miRNAs fail to be processed by Dicer and undergo degradation. The repression of let-7 expression is required for normal development and contributes to maintain the pluripotent state by preventing let-7-mediated differentiation of embryonic stem cells (PubMed: [18951094](http://www.uniprot.org/citations/18951094) target=" _blank">18951094, PubMed: [19703396](http://www.uniprot.org/citations/19703396) target=" _blank">19703396, PubMed: [22118463](http://www.uniprot.org/citations/22118463) target=" _blank">22118463, PubMed: [22898984](http://www.uniprot.org/citations/22898984) target=" _blank">22898984). Localized to the periendoplasmic reticulum area, binds to a large number of spliced mRNAs and inhibits the translation of mRNAs destined for the ER, reducing the synthesis of transmembrane proteins, ER or Golgi lumen proteins, and secretory proteins. Binds to and enhances the translation of mRNAs for several metabolic enzymes, such as PFKF, PDHA1 or SDHA, increasing glycolysis and oxidative phosphorylation. Which, with the let-7 repression may enhance tissue repair in adult tissue (By similarity).

Cellular Location

Cytoplasm. Rough endoplasmic reticulum {ECO:0000250|UniProtKB:Q8K3Y3}. Cytoplasm, P-body. Cytoplasm, Stress granule. Nucleus, nucleolus {ECO:0000250|UniProtKB:Q8K3Y3}. Note=Predominantly cytoplasmic (PubMed:22118463). In the cytoplasm, localizes to peri-endoplasmic reticulum regions and detected in the microsomal fraction derived from rough endoplasmic reticulum (RER) following subcellular fractionation May be bound to the cytosolic surface of RER on which ER-associated mRNAs are translated (By similarity). Shuttle from the nucleus to the cytoplasm requires RNA-binding (PubMed:17617744). Nucleolar localization is observed in 10-15% of the nuclei in differentiated myotubes (By similarity). {ECO:0000250|UniProtKB:Q8K3Y3, ECO:0000269|PubMed:17617744, ECO:0000269|PubMed:22118463}

Tissue Location

Expressed in embryonic stem cells, placenta and testis. Tends to be up-regulated in HER2-overexpressing breast tumors


Lin28 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

- [Flow Cytometry](#)
- [Cell Culture](#)

Lin28 Antibody - Images



Western blot analysis of Lin28 expression in NCCIT cell lysate.