

SDHA Antibody

Rabbit mAb Catalog # AP91182

Specification

SDHA Antibody - Product Information

Application Primary Accession Reactivity Clonality Other Names PGL5; SDH1; SDH2; SDHA; SDHF;	WB, IHC, FC, ICC, IP <u>P31040</u> Rat Monoclonal	
lsotype Host Calculated MW	Rabbit IgG Rabbit 72692 Da	
SDHA Antibody - Additional Information		
Purification	Affinity-chromatography	

Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human SDHA
Description	Flavoprotein (FP) subunit of succinate dehydrogenase (SDH) that is involved in complex II of the mitochondrial electron transport chain and is responsible for transferring electrons from succinate to ubiquinone.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

SDHA Antibody - Protein Information

Name SDHA

Synonyms SDH2, SDHF

Function

Flavoprotein (FP) subunit of succinate dehydrogenase (SDH) that is involved in complex II of the mitochondrial electron transport chain and is responsible for transferring electrons from succinate to ubiquinone (coenzyme Q) (PubMed:10746566, PubMed:24781757). SDH also oxidizes malate to the non-canonical enol form of oxaloacetate, enol- oxaloacetate (By similarity). Enol-oxaloacetate, which is a potent inhibitor of the succinate dehydrogenase activity, is further isomerized into keto-oxaloacetate (By similarity). Can act as a tumor suppressor (PubMed:<a href="http://www.uniprot.org/citations/20484225"



target="_blank">20484225).

Cellular Location


Mitochondrion inner membrane; Peripheral membrane protein; Matrix side

SDHA Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- <u>Flow Cytomety</u>
- <u>Cell Culture</u>

SDHA Antibody - Images



Western blot analysis of SDHA expression in (1) HeLa cell lysate; (2) Mouse brain lysate; (3) Rat brain lysate;