

**CaMKII alpha Antibody**  
Rabbit mAb  
Catalog # AP91208**Specification****CaMKII alpha Antibody - Product Information**

Application	WB, FC, IP
Primary Accession	<a href="#">O9UQM7</a>
Reactivity	Rat
Clonality	Monoclonal
<b>Other Names</b>	
Alpha CaMKII; Camk2a; CAMKA; CaMKII; CaMKIINalpha; PK2CDD; PKCCD;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	54088 Da

**CaMKII alpha Antibody - Additional Information**

Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human CaMKII alpha
Description	CaM-kinase II (CAMK2) is a prominent kinase in the central nervous system that may function in long-term potentiation and neurotransmitter release. Member of the NMDAR signaling complex in excitatory synapses it may regulate NMDAR-dependent potentiation of the AMPAR and synaptic plasticity.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

**CaMKII alpha Antibody - Protein Information****Name** CAMK2A**Synonyms** CAMKA, KIAA0968**Function**

Calcium/calmodulin-dependent protein kinase that functions autonomously after Ca(2+)/calmodulin-binding and autophosphorylation, and is involved in various processes, such as synaptic plasticity, neurotransmitter release and long-term potentiation (PubMed:<a href="http://www.uniprot.org/citations/14722083" target="\_blank">14722083</a>). Member of the NMDAR signaling complex in excitatory synapses, it regulates NMDAR-dependent potentiation of the AMPAR and therefore excitatory synaptic transmission (By similarity). Regulates dendritic

spine development (PubMed:<a href="http://www.uniprot.org/citations/28130356" target="\_blank">28130356</a>). Also regulates the migration of developing neurons (PubMed:<a href="http://www.uniprot.org/citations/29100089" target="\_blank">29100089</a>). Phosphorylates the transcription factor FOXO3 to activate its transcriptional activity (PubMed:<a href="http://www.uniprot.org/citations/23805378" target="\_blank">23805378</a>). Phosphorylates the transcription factor ETS1 in response to calcium signaling, thereby decreasing ETS1 affinity for DNA (By similarity). In response to interferon-gamma (IFN-gamma) stimulation, catalyzes phosphorylation of STAT1, stimulating the JAK- STAT signaling pathway (PubMed:<a href="http://www.uniprot.org/citations/11972023" target="\_blank">11972023</a>). In response to interferon- beta (IFN-beta) stimulation, stimulates the JAK-STAT signaling pathway (PubMed:<a href="http://www.uniprot.org/citations/35568036" target="\_blank">35568036</a>). Acts as a negative regulator of 2- arachidonoylglycerol (2-AG)-mediated synaptic signaling via modulation of DAGLA activity (By similarity).

### Cellular Location


Synapse {ECO:0000250|UniProtKB:P11275}. Postsynaptic density {ECO:0000250|UniProtKB:P11275}. Cell projection, dendritic spine. Cell projection, dendrite. Note=Postsynaptic lipid rafts {ECO:0000250|UniProtKB:P11275}

### CaMKII alpha Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### CaMKII alpha Antibody - Images



Western blot analysis of CaMKII alpha expression in SH-SY5Y cell lysate.