

PRDX6 Antibody

Rabbit mAb Catalog # AP91214

Specification

PRDX6 Antibody - Product Information

Application WB, ICC, IP
Primary Accession P30041
Reactivity Rat
Clonality Monoclonal

Other Names

Peroxiredoxin-6; Antioxidant protein 2; Liver 2D page spot 40; NSGPx; PRDX6; AOP2; KIAA;

Isotype Rabbit IgG
Host Rabbit
Calculated MW 25035 Da

PRDX6 Antibody - Additional Information

Purification Affinity-chromatography

Immunogen A synthesized peptide derived from human

PRDX6

Description Involved in redox regulation of the cell.

Can reduce H(2)O(2) and short chain

organic, fatty acid, and phospholipid hydroperoxides. May play a role in the regulation of phospholipid turnover as well as in protection against oxidative injury. Rabbit IgG in phosphate buffered saline,

pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid

freeze / thaw cycle.

PRDX6 Antibody - Protein Information

Name PRDX6

Synonyms AOP2, KIAA0106

Storage Condition and Buffer

Function

Thiol-specific peroxidase that catalyzes the reduction of hydrogen peroxide and organic hydroperoxides to water and alcohols, respectively (PubMed:10893423, PubMed:9497358). Can reduce

H(2)O(2) and short chain organic, fatty acid, and phospholipid hydroperoxides (PubMed:10893423). Also has phospholipase activity, can therefore either reduce the oxidized sn-2 fatty acyl group of phospholipids (peroxidase activity) or hydrolyze the sn-2 ester bond of phospholipids



(phospholipase activity) (PubMed:10893423, PubMed:26830860). These activities are dependent on binding to phospholipids at acidic pH and to oxidized phospholipids at cytosolic pH (PubMed:10893423). Plays a role in cell protection against oxidative stress by detoxifying peroxides and in phospholipid homeostasis (PubMed:<a href="http://www.uniprot.org/citations/10893423"

target="_blank">10893423). Exhibits acyl-CoA-dependent lysophospholipid acyltransferase which mediates the conversion of lysophosphatidylcholine (1-acyl-sn-glycero-3- phosphocholine or LPC) into phosphatidylcholine (1,2-diacyl-sn-glycero- 3-phosphocholine or PC) (PubMed:26830860). Shows a clear preference for LPC as the lysophospholipid and for palmitoyl CoA as the fatty acyl substrate (PubMed:26830860).

Cellular Location


Cytoplasm. Lysosome {ECO:0000250|UniProtKB:O35244}. Note=Also found in lung secretory organelles (lamellar bodies). {ECO:0000250|UniProtKB:O35244}

PRDX6 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

PRDX6 Antibody - Images



Western blot analysis of PRDX6 expression in HeLa cell lysate.