

SDHB Antibody

Rabbit mAb Catalog # AP91290

Specification

SDHB Antibody - Product Information

ApplicationWB, IHC, FC, IPPrimary AccessionP21912ReactivityRatClonalityMonoclonalOther NamesIron-sulfur subunit of complex II; SDHB; SDH; SDH1; PGL4;

Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	31630 Da

SDHB Antibody - Additional Information

Purification Immunogen	Affinity-chromatography A synthesized peptide derived from human SDHB
Description	Iron-sulfur protein (IP) subunit of succinate dehydrogenase (SDH) that is involved in complex II of the mitochondrial electron transport chain and is responsible for transferring electrons from succinate to ubiquinone (coenzyme Q).
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

SDHB Antibody - Protein Information

Name SDHB

Synonyms SDH, SDH1

Function

Iron-sulfur protein (IP) subunit of the succinate dehydrogenase complex (mitochondrial respiratory chain complex II), responsible for transferring electrons from succinate to ubiquinone (coenzyme Q) (PubMed:26925370, PubMed:26925370, PubMed:26925370, PubMed:26925370, PubMed:26925370, PubMed:26925370, PubMed:27604842). SDH also oxidizes malate to the non-canonical enol form of

oxaloacetate, enol- oxaloacetate (By similarity). Enol-oxaloacetate, which is a potent inhibitor of the succinate dehydrogenase activity, is further isomerized into keto-oxaloacetate (By similarity).



Cellular Location


Mitochondrion inner membrane; Peripheral membrane protein; Matrix side

SDHB Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

SDHB Antibody - Images



Western blot analysis of SDHB expression in A431 cell lysate.