

**GNAO1 Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP9134b**

**Specification**

---

**GNAO1 Antibody (C-term) - Product Information**

Application	WB, FC,E
Primary Accession	<a href="#">P09471</a>
Other Accession	<a href="#">P08239</a>
Reactivity	Mouse, Rat
Predicted	Bovine
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	291-320

**GNAO1 Antibody (C-term) - Additional Information**

**Gene ID** 2775

**Other Names**

Guanine nucleotide-binding protein G(o) subunit alpha, GNAO1

**Target/Specificity**

This GNAO1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 291-320 amino acids from the C-terminal region of human GNAO1.

**Dilution**

WB~~1:2000

FC~~1:10~50

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

GNAO1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**GNAO1 Antibody (C-term) - Protein Information**

**Name** GNAO1

**Function** Guanine nucleotide-binding proteins (G proteins) function as transducers downstream of

G protein-coupled receptors (GPCRs) in numerous signaling cascades (PubMed:[29925951](#)). The alpha chain contains the guanine nucleotide binding site and alternates between an active, GTP-bound state and an inactive, GDP-bound state (By similarity). Signaling by an activated GPCR promotes GDP release and GTP binding (By similarity). The alpha subunit has a low GTPase activity that converts bound GTP to GDP, thereby terminating the signal (By similarity). Both GDP release and GTP hydrolysis are modulated by numerous regulatory proteins (By similarity). Signaling is mediated via effector proteins, such as adenylate cyclase (By similarity). Inhibits adenylate cyclase activity, leading to decreased intracellular cAMP levels (By similarity).

#### Cellular Location


Cell membrane. Membrane; Lipid-anchor

#### GNAO1 Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### GNAO1 Antibody (C-term) - Images



All lanes : Anti-GNAO1 Antibody (C-term) at 1:2000 dilution Lane 1: Mouse cerebellum whole lysate Lane 2: Mouse brain whole lysate Lane 3: Rat brain whole lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 40 kDa Blocking/Dilution buffer: 5% NFDN/TBST.



GNAO1 Antibody (C-term) (Cat. #AP9134b) flow cytometry analysis of K562 cells (bottom histogram) compared to a negative control cell (top histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

#### **GNAO1 Antibody (C-term) - Background**

Guanine nucleotide-binding proteins (G proteins) are involved as modulators or transducers in various transmembrane signaling systems. The G(o) protein function is not clear.

#### **GNAO1 Antibody (C-term) - References**

Yi, F., et al., J. Biol. Chem. 266 (6), 3900-3906 (1991)