

FOLR1 Antibody
Rabbit mAb
Catalog # AP91419

Specification

FOLR1 Antibody - Product Information

Application	WB
Primary Accession	P15328
Reactivity	Rat
Clonality	Monoclonal

Other Names

Adult folate binding protein; FBP; Folate receptor 1; Folate Receptor 1 Precursor; Folate receptor alpha; FOLR1; FR alpha; KB cells FBP;

Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	29819 Da

FOLR1 Antibody - Additional Information

Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human FOLR1
Description	Binds to folate and reduced folic acid derivatives and mediates delivery of 5-methyltetrahydrofolate to the interior of cells.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

FOLR1 Antibody - Protein Information

Name FOLR1

Synonyms FOLR

Function

Binds to folate and reduced folic acid derivatives and mediates delivery of 5-methyltetrahydrofolate and folate analogs into the interior of cells (PubMed:[19074442](http://www.uniprot.org/citations/19074442)), PubMed:[23851396](http://www.uniprot.org/citations/23851396)), PubMed:[23934049](http://www.uniprot.org/citations/23934049)), PubMed:[2527252](http://www.uniprot.org/citations/2527252)), PubMed:[8033114](http://www.uniprot.org/citations/8033114)), PubMed:[8567728](http://www.uniprot.org/citations/8567728)). Has high affinity for folate and folic acid analogs at neutral pH (PubMed:[8567728](http://www.uniprot.org/citations/8567728)).

href="http://www.uniprot.org/citations/23851396" target="_blank">23851396, PubMed:23934049, PubMed:2527252, PubMed:8033114, PubMed:8567728). Exposure to slightly acidic pH after receptor endocytosis triggers a conformation change that strongly reduces its affinity for folates and mediates their release (PubMed:8567728). Required for normal embryonic development and normal cell proliferation (By similarity).

Cellular Location

Cell membrane; Lipid-anchor, GPI-anchor Apical cell membrane; Lipid-anchor, GPI- anchor Basolateral cell membrane; Lipid-anchor, GPI-like-anchor. Secreted Cytoplasmic vesicle. Cytoplasmic vesicle, clathrin-coated vesicle. Endosome. Note=Endocytosed into cytoplasmic vesicles and then recycled to the cell membrane

Tissue Location


Primarily expressed in tissues of epithelial origin. Expression is increased in malignant tissues. Expressed in kidney, lung and cerebellum. Detected in placenta and thymus epithelium.

FOLR1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FOLR1 Antibody - Images



Western blot analysis of FOLR1 expression in HEK293 cell lysate.