

RPL18A Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP9141b**Specification**

RPL18A Antibody (C-term) - Product Information

Application	WB, IHC-P, FC,E
Primary Accession	Q02543
Other Accession	P62718 , P62717 , Q7ZWI4 , Q3T003 , G1TTY7
Reactivity	Human, Mouse
Predicted	Bovine, Zebrafish, Rabbit, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	140-166

RPL18A Antibody (C-term) - Additional Information**Gene ID** 6142**Other Names**

60S ribosomal protein L18a, RPL18A

Target/Specificity

This RPL18A antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 140-166 amino acids from the C-terminal region of human RPL18A.

Dilution

WB~~1:1000
IHC-P~~1:50~100
FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

RPL18A Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

RPL18A Antibody (C-term) - Protein Information**Name** RPL18A

Function Component of the large ribosomal subunit. The ribosome is a large ribonucleoprotein complex responsible for the synthesis of proteins in the cell.

Cellular Location


Cytoplasm.

RPL18A Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


RPL18A Antibody (C-term) - Images




Western blot analysis of RPL18A Antibody (C-term) (Cat. #AP9141b) in 293, mouse NIH-3T3 cell line lysates (35ug/lane). RPL18A (arrow) was detected using the purified Pab.



All lanes : Anti-RPL18A Antibody (C-term) at 1:1000 dilution Lane 1: HeLa whole cell lysate Lane 2: MOLT-4 whole cell lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 21 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Formalin-fixed and paraffin-embedded human brain tissue reacted with RPL18A Antibody (C-term), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.



RPL18A Antibody (C-term) (Cat. #AP9141b) flow cytometric analysis of 293 cells (right histogram)

compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

RPL18A Antibody (C-term) - Background

Ribosomes, the organelles that catalyze protein synthesis, consist of a small 40S subunit and a large 60S subunit. Together these subunits are composed of 4 RNA species and approximately 80 structurally distinct proteins. RPL18A is a ribosomal protein that is a component of the 60S subunit. The protein belongs to the L18AE family of ribosomal proteins. It is located in the cytoplasm.

RPL18A Antibody (C-term) - References

Wool,I.G., et.al., Biochem. Cell Biol. 73 (11-12), 933-947 (1995)