

**SKP2 Antibody**  
Rabbit mAb  
Catalog # AP91547

## Specification

---

### SKP2 Antibody - Product Information

Application	WB, IHC, FC, ICC
Primary Accession	<a href="#">O13309</a>
Reactivity	Rat
Clonality	Monoclonal
<b>Other Names</b>	
FBL1; FLB1; FBXL1; MGC1366; SKP2;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	47761 Da

### SKP2 Antibody - Additional Information

Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human SKP2
Description	Substrate recognition component of a SCF (SKP1-CUL1-F-box protein) E3 ubiquitin-protein ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of target proteins involved in cell cycle progression, signal transduction and transcription.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

### SKP2 Antibody - Protein Information

**Name** SKP2

**Synonyms** FBXL1

#### Function

Substrate recognition component of a SCF (SKP1-CUL1-F-box protein) E3 ubiquitin-protein ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of target proteins involved in cell cycle progression, signal transduction and transcription (PubMed:<a href="http://www.uniprot.org/citations/11931757" target="\_blank">11931757</a>, PubMed:<a href="http://www.uniprot.org/citations/12435635" target="\_blank">12435635</a>, PubMed:<a href="http://www.uniprot.org/citations/12769844" target="\_blank">12769844</a>, PubMed:<a href="http://www.uniprot.org/citations/12769844" target="\_blank">12769844</a>)

href="http://www.uniprot.org/citations/12840033" target="\_blank">12840033</a>, PubMed:<a href="http://www.uniprot.org/citations/15342634" target="\_blank">15342634</a>, PubMed:<a href="http://www.uniprot.org/citations/15668399" target="\_blank">15668399</a>, PubMed:<a href="http://www.uniprot.org/citations/15949444" target="\_blank">15949444</a>, PubMed:<a href="http://www.uniprot.org/citations/16103164" target="\_blank">16103164</a>, PubMed:<a href="http://www.uniprot.org/citations/16262255" target="\_blank">16262255</a>, PubMed:<a href="http://www.uniprot.org/citations/16581786" target="\_blank">16581786</a>, PubMed:<a href="http://www.uniprot.org/citations/16951159" target="\_blank">16951159</a>, PubMed:<a href="http://www.uniprot.org/citations/17908926" target="\_blank">17908926</a>, PubMed:<a href="http://www.uniprot.org/citations/17962192" target="\_blank">17962192</a>, PubMed:<a href="http://www.uniprot.org/citations/22464731" target="\_blank">22464731</a>, PubMed:<a href="http://www.uniprot.org/citations/22770219" target="\_blank">22770219</a>, PubMed:<a href="http://www.uniprot.org/citations/32267835" target="\_blank">32267835</a>). Specifically recognizes phosphorylated CDKN1B/p27kip and is involved in regulation of G1/S transition (By similarity). Degradation of CDKN1B/p27kip also requires CKS1 (By similarity). Recognizes target proteins ORC1, CDT1, RBL2, KMT2A/MLL1, CDK9, RAG2, NBN, FOXO1, UBP43, YTHDF2, and probably MYC, TOB1 and TAL1 (PubMed:<a href="http://www.uniprot.org/citations/11931757" target="\_blank">11931757</a>, PubMed:<a href="http://www.uniprot.org/citations/12435635" target="\_blank">12435635</a>, PubMed:<a href="http://www.uniprot.org/citations/12769844" target="\_blank">12769844</a>, PubMed:<a href="http://www.uniprot.org/citations/12840033" target="\_blank">12840033</a>, PubMed:<a href="http://www.uniprot.org/citations/15342634" target="\_blank">15342634</a>, PubMed:<a href="http://www.uniprot.org/citations/15668399" target="\_blank">15668399</a>, PubMed:<a href="http://www.uniprot.org/citations/15949444" target="\_blank">15949444</a>, PubMed:<a href="http://www.uniprot.org/citations/16103164" target="\_blank">16103164</a>, PubMed:<a href="http://www.uniprot.org/citations/16581786" target="\_blank">16581786</a>, PubMed:<a href="http://www.uniprot.org/citations/16951159" target="\_blank">16951159</a>, PubMed:<a href="http://www.uniprot.org/citations/17908926" target="\_blank">17908926</a>, PubMed:<a href="http://www.uniprot.org/citations/17962192" target="\_blank">17962192</a>, PubMed:<a href="http://www.uniprot.org/citations/22464731" target="\_blank">22464731</a>, PubMed:<a href="http://www.uniprot.org/citations/32267835" target="\_blank">32267835</a>). Degradation of TAL1 also requires STUB1 (PubMed:<a href="http://www.uniprot.org/citations/17962192" target="\_blank">17962192</a>). Recognizes CDKN1A in association with CCNE1 or CCNE2 and CDK2 (PubMed:<a href="http://www.uniprot.org/citations/16262255" target="\_blank">16262255</a>). Promotes ubiquitination and destruction of CDH1 in a CK1-dependent manner, thereby regulating cell migration (PubMed:<a href="http://www.uniprot.org/citations/22770219" target="\_blank">22770219</a>). Following phosphorylation in response to DNA damage, mediates 'Lys-63'-linked ubiquitination of NBN, promoting ATM recruitment to DNA damage sites and DNA repair via homologous recombination (PubMed:<a href="http://www.uniprot.org/citations/22464731" target="\_blank">22464731</a>).

### Cellular Location

Cytoplasm. Nucleus


### SKP2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)

- [Cell Culture](#)

## SKP2 Antibody - Images



Western blot analysis of SKP2 expression in (1) Jurkat cell lysate; (2) NIH/3T3 cell lysate.