

FADD Antibody
Rabbit mAb
Catalog # AP91555

Specification

FADD Antibody - Product Information

Application	WB, IHC, FC, IP
Primary Accession	Q13158
Clonality	Monoclonal
Other Names	
FADD protein; MORT1;	

Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	23279 Da

FADD Antibody - Additional Information

Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human FADD
Description	Apoptotic adaptor molecule that recruits caspase-8 or caspase-10 to the activated Fas (CD95) or TNFR-1 receptors. The resulting aggregate called the death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation. Active caspase-8 initiates the subsequent cascade of caspases mediating apoptosis. Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
Storage Condition and Buffer	

FADD Antibody - Protein Information

Name FADD {ECO:0000303|PubMed:7538907, ECO:0000312|HGNC:HGNC:3573}

Function

Apoptotic adapter molecule that recruits caspases CASP8 or CASP10 to the activated FAS/CD95 or TNFRSF1A/TNFR-1 receptors (PubMed: [16762833](http://www.uniprot.org/citations/16762833), PubMed: [19118384](http://www.uniprot.org/citations/19118384), PubMed: [20935634](http://www.uniprot.org/citations/20935634), PubMed: [23955153](http://www.uniprot.org/citations/23955153), PubMed: [24025841](http://www.uniprot.org/citations/24025841), PubMed: [7538907](http://www.uniprot.org/citations/7538907), PubMed: [9184224](http://www.uniprot.org/citations/9184224)). The resulting aggregate called the death-inducing signaling

complex (DISC) performs CASP8 proteolytic activation (PubMed:16762833, PubMed:19118384, PubMed:20935634, PubMed:7538907, PubMed:9184224). Active CASP8 initiates the subsequent cascade of caspases mediating apoptosis (PubMed:16762833). Involved in interferon-mediated antiviral immune response, playing a role in the positive regulation of interferon signaling (PubMed:21109225).

Tissue Location


Expressed in a wide variety of tissues, except for peripheral blood mononuclear leukocytes.

FADD Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FADD Antibody - Images



Western blot analysis of FADD expression in A431 cell lysate.