

Mad2L2 Antibody
Rabbit mAb
Catalog # AP91576

Specification

Mad2L2 Antibody - Product Information

Application	WB, IHC, FC, ICC, IP
Primary Accession	O9UI95
Reactivity	Rat
Clonality	Monoclonal
Other Names	
hREV7; MAD2B; Mad2L2 Mitotic Arrest Deficient 2 L2; POLZ2; REV 7; REV7; REV7 homolog;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	24334 Da

Mad2L2 Antibody - Additional Information

Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Mad2L2
Description	Adapter protein able to interact with different proteins and involved in different biological processes. Mediates the interaction between the error-prone DNA polymerase zeta catalytic subunit REV3L and the inserter polymerase REV1, thereby mediating the second polymerase switching in translesion DNA synthesis.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Mad2L2 Antibody - Protein Information

Name MAD2L2

Synonyms MAD2B, REV7

Function

Adapter protein able to interact with different proteins and involved in different biological processes (PubMed: [11459825](http://www.uniprot.org/citations/11459825) target="_blank">11459825, PubMed: [11459826](http://www.uniprot.org/citations/11459826) target="_blank">11459826, PubMed: [17296730](http://www.uniprot.org/citations/17296730) target="_blank">17296730, PubMed: [17719540](http://www.uniprot.org/citations/17719540) target="_blank">17719540, PubMed: [19443654](http://www.uniprot.org/citations/19443654) target="_blank">19443654)

target="_blank">19443654, PubMed:29656893). Mediates the interaction between the error-prone DNA polymerase zeta catalytic subunit REV3L and the inserter polymerase REV1, thereby mediating the second polymerase switching in translesion DNA synthesis (PubMed:20164194). Translesion DNA synthesis releases the replication blockade of replicative polymerases, stalled in presence of DNA lesions (PubMed:20164194). Component of the shieldin complex, which plays an important role in repair of DNA double-stranded breaks (DSBs) (PubMed:29656893). During G1 and S phase of the cell cycle, the complex functions downstream of TP53BP1 to promote non-homologous end joining (NHEJ) and suppress DNA end resection (PubMed:29656893). Mediates various NHEJ-dependent processes including immunoglobulin class-switch recombination, and fusion of unprotected telomeres (PubMed:29656893). May also regulate another aspect of cellular response to DNA damage through regulation of the JNK-mediated phosphorylation and activation of the transcriptional activator ELK1 (PubMed:17296730). Inhibits the FZR1- and probably CDC20-mediated activation of the anaphase promoting complex APC thereby regulating progression through the cell cycle (PubMed:11459825, PubMed:17719540). Regulates TCF7L2-mediated gene transcription and may play a role in epithelial-mesenchymal transdifferentiation (PubMed:19443654).

Cellular Location

Nucleus. Cytoplasm, cytoskeleton, spindle. Cytoplasm. Chromosome. Note=Recruited to sites of chromosomal double-stranded breaks during G1 and S phase of the cell cycle

Tissue Location

Ubiquitously expressed.


Mad2L2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Mad2L2 Antibody - Images





Western blot analysis of Mad2L2 expression in HeLa cell lysate.