

**TRIM25 Antibody**  
Rabbit mAb  
Catalog # AP91584

## Specification

---

### TRIM25 Antibody - Product Information

Application	WB, IHC, FC, ICC, IP
Primary Accession	<a href="#">Q14258</a>
Reactivity	Rat
Clonality	Monoclonal
<b>Other Names</b>	
EFP; RNF147; TRI25; TRIM 25; Trim25; Z147; ZNF147;	

Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	70973 Da

### TRIM25 Antibody - Additional Information

Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human TRIM25
Description	Functions as an ubiquitin E3 ligase and as an ISG15 E3 ligase. Involved in innate immune defense against viruses by mediating ubiquitination of DDX58. Mediates 'Lys-63'-linked polyubiquitination of the DDX58 N-terminal CARD-like region which is crucial for triggering the cytosolic signal transduction that leads to the production of interferons in response to viral infection.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

### TRIM25 Antibody - Protein Information

**Name** TRIM25

**Synonyms** EFP {ECO:0000303|PubMed:8248217}, RNF147

#### Function

Functions as a ubiquitin E3 ligase and as an ISG15 E3 ligase (PubMed:<a href="http://www.uniprot.org/citations/16352599" target="\_blank">16352599</a>). Involved in innate immune defense against viruses by mediating ubiquitination of RIGI and IFIH1 (PubMed:<a href="http://www.uniprot.org/citations/17392790" target="\_blank">17392790</a>, PubMed:<a href="http://www.uniprot.org/citations/17392790" target="\_blank">17392790</a>).

<http://www.uniprot.org/citations/29357390> target="\_blank">29357390</a>, PubMed:<a href="http://www.uniprot.org/citations/30193849" target="\_blank">30193849</a>, PubMed:<a href="http://www.uniprot.org/citations/31710640" target="\_blank">31710640</a>, PubMed:<a href="http://www.uniprot.org/citations/33849980" target="\_blank">33849980</a>, PubMed:<a href="http://www.uniprot.org/citations/36045682" target="\_blank">36045682</a>). Mediates 'Lys-63'-linked polyubiquitination of the RIGI N-terminal CARD-like region and may play a role in signal transduction that leads to the production of interferons in response to viral infection (PubMed:<a href="http://www.uniprot.org/citations/17392790" target="\_blank">17392790</a>, PubMed:<a href="http://www.uniprot.org/citations/23950712" target="\_blank">23950712</a>). Mediates 'Lys-63'-linked polyubiquitination of IFIH1 (PubMed:<a href="http://www.uniprot.org/citations/30193849" target="\_blank">30193849</a>). Promotes ISGylation of 14-3-3 sigma (SFN), an adapter protein implicated in the regulation of a large spectrum signaling pathway (PubMed:<a href="http://www.uniprot.org/citations/16352599" target="\_blank">16352599</a>, PubMed:<a href="http://www.uniprot.org/citations/17069755" target="\_blank">17069755</a>). Mediates estrogen action in various target organs (PubMed:<a href="http://www.uniprot.org/citations/22452784" target="\_blank">22452784</a>). Mediates the ubiquitination and subsequent proteasomal degradation of ZFHX3 (PubMed:<a href="http://www.uniprot.org/citations/22452784" target="\_blank">22452784</a>). Plays a role in promoting the restart of stalled replication forks via interaction with the KHDC3L-OOEP scaffold and subsequent ubiquitination of BLM, resulting in the recruitment and retainment of BLM at DNA replication forks (By similarity). Plays an essential role in the antiviral activity of ZAP/ZC3HAV1; an antiviral protein which inhibits the replication of certain viruses. Mechanistically, mediates 'Lys-63'-linked polyubiquitination of ZAP/ZC3HAV1 that is required for its optimal binding to target mRNA (PubMed:<a href="http://www.uniprot.org/citations/28060952" target="\_blank">28060952</a>, PubMed:<a href="http://www.uniprot.org/citations/28202764" target="\_blank">28202764</a>). Mediates also the ubiquitination of various substrates implicated in stress granule formation, nonsense-mediated mRNA decay, nucleoside synthesis and mRNA translation and stability (PubMed:<a href="http://www.uniprot.org/citations/36067236" target="\_blank">36067236</a>).

#### Cellular Location

Cytoplasm. Cytoplasm, Stress granule. Nucleus {ECO:0000250|UniProtKB:Q61510}

#### Tissue Location

Expressed in breast tumors (at protein level). Ubiquitous.


### TRIM25 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### TRIM25 Antibody - Images





Western blot analysis of TRIM25 expression in HeLa cell lysate.