

**JAK1 Antibody**  
Rabbit mAb  
Catalog # AP91603

## Specification

---

### JAK1 Antibody - Product Information

Application	WB
Primary Accession	<a href="#">P23458</a>
Reactivity	Rat
Clonality	Monoclonal
<b>Other Names</b>	
JAK 1; JAK 1A; JAK 1B; JAK1; JAK1A; JAK1B; JTK3;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	133277 Da

### JAK1 Antibody - Additional Information

Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human JAK1
Description	Tyrosine kinase of the non-receptor type, involved in the IFN-alpha/beta/gamma signal pathway. Kinase partner for the interleukin (IL)-2 receptor.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

### JAK1 Antibody - Protein Information

**Name** JAK1

**Synonyms** JAK1A, JAK1B

#### Function

Tyrosine kinase of the non-receptor type, involved in the IFN-alpha/beta/gamma signal pathway (PubMed: [16239216](http://www.uniprot.org/citations/16239216)), (PubMed: [28111307](http://www.uniprot.org/citations/28111307)), (PubMed: [32750333](http://www.uniprot.org/citations/32750333)), (PubMed: [7615558](http://www.uniprot.org/citations/7615558)), (PubMed: [8232552](http://www.uniprot.org/citations/8232552)). Kinase partner for the interleukin (IL)-2 receptor (PubMed: [11909529](http://www.uniprot.org/citations/11909529)) as well as interleukin (IL)-10 receptor (PubMed: [12133952](http://www.uniprot.org/citations/12133952)). Kinase partner for the type I interferon receptor IFNAR2

(PubMed:<a href="http://www.uniprot.org/citations/16239216" target="\_blank">16239216</a>, PubMed:<a href="http://www.uniprot.org/citations/28111307" target="\_blank">28111307</a>, PubMed:<a href="http://www.uniprot.org/citations/32750333" target="\_blank">32750333</a>, PubMed:<a href="http://www.uniprot.org/citations/7615558" target="\_blank">7615558</a>, PubMed:<a href="http://www.uniprot.org/citations/8232552" target="\_blank">8232552</a>). In response to interferon-binding to IFNAR1-IFNAR2 heterodimer, phosphorylates and activates its binding partner IFNAR2, creating docking sites for STAT proteins (PubMed:<a href="http://www.uniprot.org/citations/7759950" target="\_blank">7759950</a>). Directly phosphorylates STAT proteins but also activates STAT signaling through the transactivation of other JAK kinases associated with signaling receptors (PubMed:<a href="http://www.uniprot.org/citations/16239216" target="\_blank">16239216</a>, PubMed:<a href="http://www.uniprot.org/citations/32750333" target="\_blank">32750333</a>, PubMed:<a href="http://www.uniprot.org/citations/8232552" target="\_blank">8232552</a>).

### Cellular Location

Endomembrane system; Peripheral membrane protein. Note=Wholly intracellular, possibly membrane associated

### Tissue Location


Expressed at higher levels in primary colon tumors than in normal colon tissue. The expression level in metastatic colon tumors is comparable to the expression level in normal colon tissue

## JAK1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## JAK1 Antibody - Images



Western blot analysis of JAK1 expression in (1) A431 cell lysate; (2) RAW264.7 cell lysate.