

Egr1 Antibody
Rabbit mAb
Catalog # AP91607

Specification

Egr1 Antibody - Product Information

Application	WB, IHC, ICC
Primary Accession	P18146
Clonality	Monoclonal
Other Names	
TIS8; AT225; G0S30; NGFI-A; ZNF225; KROX-24; ZIF-268; EGR1;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	57507 Da

Egr1 Antibody - Additional Information

Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human Egr1
Description	Transcriptional regulator. Recognizes and binds to the DNA sequence 5'-CGCCCCCGC-3'(EGR-site). Activates the transcription of target genes whose products are required for mitogenesis and differentiation.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

Egr1 Antibody - Protein Information

Name EGR1

Synonyms KROX24, ZNF225 {ECO:0000303|PubMed:21103}

Function

Transcriptional regulator (PubMed: 20121949). Recognizes and binds to the DNA sequence 5'-GCG(T/G)GGGCG-3'(EGR-site) in the promoter region of target genes (By similarity). Binds double-stranded target DNA, irrespective of the cytosine methylation status (PubMed: 25258363, PubMed: 25999311). Regulates the transcription of numerous target genes, and thereby plays an important role in regulating the response to growth factors, DNA damage, and ischemia. Plays a role in the regulation of cell survival, proliferation and cell death. Activates expression of p53/TP53 and TGFB1, and thereby

helps prevent tumor formation. Required for normal progress through mitosis and normal proliferation of hepatocytes after partial hepatectomy. Mediates responses to ischemia and hypoxia; regulates the expression of proteins such as IL1B and CXCL2 that are involved in inflammatory processes and development of tissue damage after ischemia. Regulates biosynthesis of luteinizing hormone (LHB) in the pituitary (By similarity). Regulates the amplitude of the expression rhythms of clock genes: BMAL1, PER2 and NR1D1 in the liver via the activation of PER1 (clock repressor) transcription. Regulates the rhythmic expression of core-clock gene BMAL1 in the suprachiasmatic nucleus (SCN) (By similarity).

Cellular Location

Nucleus. Cytoplasm

Tissue Location


Detected in neutrophils (at protein level).

Egr1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Egr1 Antibody - Images



Western blot analysis of Egr1 expression in 293T cell lysate treated with 20% FBS.