

SMARCC1/BAF155 Antibody
Rabbit mAb
Catalog # AP91728**Specification**

SMARCC1/BAF155 Antibody - Product Information

Application	WB, FC, ICC, IP
Primary Accession	O92922
Reactivity	Rat
Clonality	Monoclonal
Other Names	
BAF155; CRACC1; Rsc8; SMARCC1; SRG3; SWI3;	
Isotype	Rabbit IgG
Host	Rabbit
Calculated MW	122867 Da

SMARCC1/BAF155 Antibody - Additional Information

Purification	Affinity-chromatography
Immunogen	A synthesized peptide derived from human SMARCC1/BAF155
Description	Involved in transcriptional activation and repression of select genes by chromatin remodeling (alteration of DNA-nucleosome topology). May stimulate the ATPase activity of the catalytic subunit of the complex.
Storage Condition and Buffer	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.

SMARCC1/BAF155 Antibody - Protein InformationName SMARCC1 ([HGNC:11104](#))

Synonyms BAF155

Function

Involved in transcriptional activation and repression of select genes by chromatin remodeling (alteration of DNA-nucleosome topology). Component of SWI/SNF chromatin remodeling complexes that carry out key enzymatic activities, changing chromatin structure by altering DNA-histone contacts within a nucleosome in an ATP-dependent manner. May stimulate the ATPase activity of the catalytic subunit of the complex (PubMed:[10078207](http://www.uniprot.org/citations/10078207), PubMed:[29374058](http://www.uniprot.org/citations/29374058)). Belongs to the neural progenitors-specific chromatin remodeling complex (npBAF complex) and the

neuron-specific chromatin remodeling complex (nBAF complex). During neural development a switch from a stem/progenitor to a postmitotic chromatin remodeling mechanism occurs as neurons exit the cell cycle and become committed to their adult state. The transition from proliferating neural stem/progenitor cells to postmitotic neurons requires a switch in subunit composition of the npBAF and nBAF complexes. As neural progenitors exit mitosis and differentiate into neurons, npBAF complexes which contain ACTL6A/BAF53A and PHF10/BAF45A, are exchanged for homologous alternative ACTL6B/BAF53B and DPF1/BAF45B or DPF3/BAF45C subunits in neuron-specific complexes (nBAF). The npBAF complex is essential for the self-renewal/proliferative capacity of the multipotent neural stem cells. The nBAF complex along with CREST plays a role regulating the activity of genes essential for dendrite growth (By similarity).

Cellular Location

Nucleus. Cytoplasm

Tissue Location


Expressed in brain, heart, muscle, placenta, lung, liver, muscle, kidney and pancreas

SMARCC1/BAF155 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SMARCC1/BAF155 Antibody - Images



Western blot analysis of SMARCC1 expression in HeLa cell lysate.